



REC'D 24 OCT 2003

WIPO PCT

# Kongeriget Danmark

Patent application No.: PA 2003 00600

Date of filing: 15 April 2003

Applicant: Pantheco A/S  
(Name and address) Bøge Allé 3  
DK-2970 Hørsholm  
Denmark

Title: Modified PNA Molecules

IPC: -

This is to certify that the attached documents are exact copies of the above mentioned patent application as originally filed.

**PRIORITY  
DOCUMENT**  
SUBMITTED OR TRANSMITTED IN  
COMPLIANCE WITH RULE 17.1(a) OR (b)

Patent- og Varemærkestyrelsen  
Økonomi- og Erhvervsministeriet

14 October 2003

  
Pia Høybye-Olsen



15 APR. 2003

PVS

## MODIFIED PNA MOLECULES

The present invention concerns novel drugs for use in combating various diseases. More particular the invention concerns peptide nucleic acid (PNA) drugs, which are optionally  
5 modified in order to obtain novel PNA molecules with cell-specific delivery.

## BACKGROUND OF THE INVENTION

10 Antisense agents offer a novel strategy in combating diseases, as well as opportunities to employ new chemical classes in the drug design.

Oligonucleotides can interact with native DNA and RNA in several ways. One of these is duplex formation between an oligonucleotide and a single stranded nucleic acid. Another is tri-  
15 plex formation between an oligonucleotide and double stranded DNA to form a triplex structure.

Results from basic research have been encouraging, and antisense oligonucleotide drug formulations against viral and disease causing human genes are progressing through clinical  
20 trials. Efficient antisense inhibition of bacterial genes also has wide applications.

Peptide nucleic acids (PNA) are compounds that in certain respects are similar to oligonucleotides and their analogs and thus may mimic DNA and RNA. In PNA, the deoxyribose backbone of oligonucleotides has been replaced by a pseudo-peptide backbone (Nielsen et al. 1991 (1)). Each subunit, or monomer, has a naturally occurring or non-naturally occurring  
25 nucleobase attached to this backbone. One such backbone is constructed of repeating units of N-(2-aminoethyl)glycine linked through amide bonds. PNA hybridises with complementary nucleic acids through Watson and Crick base pairing and helix formation (Egholm et al. 1993 (2)). The Pseudo-peptide backbone provides superior hybridization properties (Egholm et al. 1993 (2)), resistance to enzymatic degradation (Demidov et al. 1994 (3)) and access to a variety of chemical modifications (Nielsen and Haaima 1997 (4), WO 94/25472, WO98/03542).  
30

PNA binds both DNA and RNA to form PNA/DNA or PNA/RNA duplexes. The resulting PNA/DNA or PNA/RNA duplexes are bound with greater affinity than corresponding  
35 DNA/DNA or DNA/RNA duplexes as determined by the melting point temperature ( $T_m$ ). This high thermal stability might be attributed to the lack of charge repulsion due to the neutral

backbone in PNA. In addition to increased affinity, PNA has also been shown to bind to DNA with increased specificity. When a PNA/DNA duplex mismatch is melted relative to the DNA/DNA duplex, there is seen an 8 °C to 20°C drop in the melting point temperature.

- 5 Furthermore, homopyrimidine PNA oligomers form extremely stable PNA<sub>2</sub>-DNA (RNA) triplexes with sequence complementary targets in DNA or RNA oligomers. Finally, PNA's may bind to double stranded DNA or RNA by helix invasion.

10 An advantage of PNA compared to oligonucleotides is that the PNA polyamide backbone (having appropriate nucleobases or other side chain groups attached thereto) is not recognised by either nucleases or proteases and are thus not cleaved. As a result, PNA's are resistant to degradation by enzymes unlike nucleic acids and peptides.

15 For antigenic and antisense application, target bound PNA can cause steric hindrance of DNA and RNA polymerases, reverse transcription, telomerase and of the ribosomes (Harvey et al. 1992 (5), Knudsen et al. 1996 (6), Good and Nielsen 1998 (7,8)), by targeting, among others, DNA, mRNA, rRNA, or tRNA.

20 A general difficulty when using antisense agents is cell uptake and targeting of specific organs. A variety of strategies to improve uptake can be envisioned and there are reports of improved uptake into eukaryotic cells using lipids (Lewis et al. 1996 (9)), encapsulation (Meyer et al. 1998 (10)) and carrier strategies (Nyce and Metzger 1997 (11), Pooga et al, 1998 (12)).

25 WO 99/05302 discloses a PNA conjugate consisting of PNA and the transporter peptide transportan, which peptide may be used for transport across a lipid membrane and for delivery of the PNA into interactive contact with intracellular polynucleotides.

30 US-A-5 777 078 discloses a pore-forming compound, which comprises a delivery agent recognising the target cell and being linked to a pore-forming agent, such as a bacterial exotoxin. The compound is administered together with a drug such as PNA.

35 WO 96/11205 discloses PNA conjugates, wherein a conjugated moiety may be placed on terminal or non-terminal parts of the backbone of PNA in order to functionalise the PNA. The conjugated moieties may be reporter enzymes or molecules, steroids, carbohydrate, ter-

penes, peptides, proteins, etc. It is suggested that the conjugates among other properties may possess improved transfer properties for crossing cellular membranes.

WO 01/27261 discloses conjugates of cationic peptides and aeg-PNA (cf. Figure 4).

5

WO 98/52614 discloses a method of enhancing transport over biological membranes. According to this publication, biological active agents such as PNA may be conjugated to a transporter polymer in order to enhance the transmembrane transport. The transporter polymer consists of 6-25 subunits; at least 50% of which contain a guanidino or amidino sidechain moiety and wherein at least 6 contiguous subunits contain guanidino and/or amidino sidechains. A preferred transporter polymer is a polypeptide containing nine arginine subunits ((Arg)<sub>9</sub>).

10

However, the present methods of transport of PNA oligomers across biological membranes lack efficiency and specificity. Only little information is available on the pharmacokinetic behaviour of PNA oligomers, e.g. the dynamic and kinetic mechanisms of exogenous absorption, biotransformation, distribution, release, transport, uptake, and elimination of PNA oligomers as a function of dosage and extent and rate of metabolic processes. However, data indicates that PNA oligomers are fairly quickly excreted in the urine according to McMahon et al (2002 (13)), being a rather hydrophilic compound, which rarely binds to proteins like albumin in serum.

20

It could be of significant medicinal interest to functionalise the PNAs, in order to control the bio-distribution of the molecule. Functionalisation of the PNA backbone may dramatically change the physico-chemical properties of the PNA, and it is plausible that such changes would significantly influence its pharmacokinetic behaviour.

25

Zhang et al (2001 (14)) describes a method by which uptake of PNA oligomers in a liver cell was promoted by modifying the terminal ends of PNA with lactose. Lactose, being recognized by the hepatic asialoglycoprotein receptor, provided an efficient entry of lactose modified PNAs into HepG2 cells.

30

Biessen et al (2002, (15)) presents work in which the parenchymal liver cell uptake of antisense PNA drug was improved by targeting to the asialoglycoprotein receptor of the liver cell, a glycoconjugated antisense PNA. The PNA was conjugated with N-acetyl-galactos-

35

aminyls at the 5' end through lysine side chain linkers.


Prior art shows an increased organ and cell specificity of PNAs by end glycosylation of the PNA. However, incorporation of the saccharides into the PNA backbone would improve the medicinal chemistry opportunities, the biostability and biodistribution of the drug, resulting in lower dosage and reduced side effects.

## SUMMARY OF THE INVENTION

The present invention relates to glycosylated peptide nucleic acid (PNA) monomers. More particular, the invention concerns the incorporation of glycosylated monomers into an antisense PNA oligomer, in order to improve the cell and/or organ-specific uptake of PNAs and thereby the pharmacokinetic behavior.

It has been found that by integrating saccharides within PNA monomers and by subsequent incorporation of at least one modified PNA monomer in an antisense PNA oligomer, an enhanced efficacy and organ-specificity is observed, without major influence on the DNA or RNA hybridisation potency of the PNA. The important feature of the modified PNA molecules is the incorporation of at least one glycosylated PNA monomer in the PNA oligomer chain. This results in versatility in synthesis and design in terms of character, position and number of saccharides, which is crucial for biological activity and also high bio-stability.


Thus, the present invention concerns a PNA monomer of formula (I):



wherein B is a naturally-occurring nucleobase preferably A, T, G, or C, or a non-naturally-occurring nucleobase;


(Pr) is hydrogen or a protection group;

$R^1$ ,  $R^2$  and  $R^3$  are, independently, hydrogen, an amino acid side chain, or an  $C_{2-6}$ -alkyl, aryl, aralkyl, heteroaryl, hydroxy,  $C_{1-6}$ -alkoxy,  $C_{1-6}$ -alkylthio, hydroxy- or alkoxy- or alkylthio-substituted  $C_{1-6}$ -alkyl,  $-NR^4R^5$ , (wherein  $R^4$  and  $R^5$  independently are hydrogen,  $C_{1-6}$ -alkyl, hydroxy- or alkoxy- or alkylthio-substituted  $C_{1-6}$ -alkyl), or  $Z^1-Z^2$ , wherein  $Z^1$  is a bond or one of the radicals of formula (II):




wherein  $n$  is from 0 to 8;

and  $Z^2$  is  $\alpha$ - or  $\beta$ -forms of a monosaccharide, a disaccharide, a polysaccharide, or one of the radicals of formula (III):




provided that at least one of  $R^1$ ,  $R^2$ , or  $R^3$  is  $Z^1$ - $Z^2$ .

Enclosed is also a compound according to claim 1, wherein  $Z^1$  is one of the radicals of formula (IV):



5

Enclosed is further a compound according to claim 1 or 3, wherein  $Z^1$ - $Z^2$  comprises ligands of formula (V):



10

or beta-forms hereof.

15

Enclosed is also a peptide nucleic acid oligomer with from 4 to 50 monomers selected from the group consisting of PNA monomers and at least one monomer of claim 1, said PNA oligomer conjugated either directly or through a linking moiety to hydrogen or a reporter enzyme, a reporter molecule, a steroid, a carbohydrate, a terpene, a peptide, a protein, an aromatic lipophilic molecule, a non aromatic lipophilic molecule, a phospholipid, an intercalator, a cell receptor binding molecule, a crosslinking agent, a water soluble vitamin, a lipid

soluble vitamin, an RNA/DNA cleaving complex, a metal chelator, a porphyrin, an alkylator, or a polymeric compound selected from polymeric amines, polymeric glycols and polyethers.

5 The present invention further comprises a peptide nucleic acid molecule comprising a peptide nucleic acid oligomer with from 4 to 50  $\alpha$ -PNA monomers and one or more conjugates bound to said peptide nucleic acid either directly or through one or more linking moieties, wherein said conjugate is a reporter enzyme, a reporter molecule, a steroid, a carbohydrate, a terpene, a peptide, a protein, an aromatic lipophilic molecule, a non aromatic lipophilic molecule, a phospholipid, an intercalator, a cell receptor binding molecule, a crosslinking  
10 agent, a water soluble vitamin, a lipid soluble vitamin, an RNA/DNA cleaving complex, a metal chelator, a porphyrin, an alkylator, or a polymeric compound selected from polymeric amines, polymeric glycols and polyethers.

15 In a preferred embodiment the glycoside residues (sugar residues) are chosen from monosaccharides having a high intrinsic affinity for the asialoglycoprotein receptor, including  $\beta$ -D-galactosyl, 2-acetamido-2-deoxy-galactopyranosyl, 1-phenyl- $\beta$ -D-galactosyl, 1-propyl- $\beta$ -D-galactosyl or 1-butyl- $\beta$ -D-galactosyl.

20 In another preferred embodiment of the invention the linking moiety is an amino acid sequence of from 1 to 10 positively charged amino acids or amino acid analogues.

By the term "positively charged amino acids or amino acid analogues" is to be understood any natural or non-natural occurring amino acid or amino acid analogue which have a positive charge at physiological pH.

25 Among the positively charged amino acids and amino acid analogs may be mentioned lysine (Lys, K), arginine (Arg, R), diamino butyric acid (DAB) and ornithine (Orn). The skilled person will be aware of further positively charged amino acids and amino acid analogs.

30 In one aspect of the invention, the modified PNA molecules are used in the manufacture of medicaments for the treatment or prevention of a disease selected from bacterial and viral infections, cardiac or vascular diseases, metabolic diseases or immunological disorders or for disinfecting non-living objects.

35 In a further aspect, the invention concerns a composition for treating or preventing disease



selected from bacterial and viral infections, cardiac or vascular diseases, metabolic diseases or immunological disorders or for disinfecting non-living objects.

In another aspect, the invention concerns the treatment or prevention of disease selected from bacterial and viral infections, cardiac or vascular diseases, metabolic diseases or immunological disorders or for disinfecting non-living objects.

Preferred targeting is organ related diseases e.g. liver diseases such as hepatitis and liver cancer, known for a person skilled in the art.

Liver cancer is the fifth most common cancer worldwide. More than 400,000 cases were reported in 1990. Hepatocellular carcinoma (HCC) accounts for 80% of all liver cancer. Liver cancer can result from both viral infection and chemical exposure. Known risk factors include hepatitis B and C virus infection and exposure to aflatoxin 1. It is not known whether distinct routes to liver cancer affect the same or different cellular pathways. No mutational model has yet been developed for liver cancer as it has been for other cancers such as colon cancer.

According to the invention one of numerous described high affinity ligands for the  $\alpha v\beta 3$  integrin receptor is used for PNA delivery to angiogenic blood vessels in the treatment of diseases that depend on angiogenesis and vascular remodelling, including cancer, see for example Hood JD *et al.* Science (2002) 296(5577):2404-7.

In yet a further aspect, the present invention concerns a method of identifying specific advantageous antisense PNA sequences, which may be used in the modified PNA molecule according to the invention.

#### DETAILED DESCRIPTION OF THE INVENTION

Antisense PNAs can inhibit bacterial gene expression with gene and sequence specificity (Good and Nielsen 1998a,b (12, 13) and WO 99/13893). The approach may prove practical as a tool for functional genomics and as a source for novel antimicrobial drugs. However, improvements on standard PNA are required to increase antisense potencies. The major limit to activity appears to be cellular entry and cellular specificity. Cell membranes effectively ex-

clude the entry of large molecular weight foreign compounds, and previous results for in vitro and cellular assays seem to show that the cell barrier restricts antisense effects. Accordingly, the present invention concerns strategies to improve the activity and specificity of antisense potencies.

5

Without being bound by theory, it is believed that glycosylation of PNA oligomers lead to an improved cell specific PNA uptake. It is believed that the glycosylated peptides are recognised by receptors in cell membranes such as the hepatic asialoglycoprotein receptor thereby taken up through the glucose pathway, allowing the modified PNA molecule to cross  
10 the cell wall, reaching structures inside the cell, such as the genome, mRNA's, the ribosome, etc.

15

According to the invention, PNA molecules modified with saccharides enable specific and efficient inhibition of genes with nanomolar concentrations. Antisense potencies in this concentration are consistent with practical applications of the technology. It is believed that the present invention for the first time demonstrates that peptides with a certain pattern of glycosylation can be used as carriers to deliver PNAs across cell membranes. Further, the present invention has made it possible to administer PNA in an efficient concentration, which is also acceptable to the patient.

20

The terms "C<sub>1-6</sub>-alkyl" as used herein, represent a branched or straight alkyl group having from one to six carbon atoms. Typical C<sub>1-6</sub>-alkyl groups include, but are not limited to, methyl, ethyl, n-propyl, iso-propyl, butyl, iso-butyl, sec-butyl, tert-butyl, pentyl, iso-pentyl, hexyl, iso-hexyl and the like.

25

The number of modified PNA monomers in the PNA oligomer may be chosen between 1 and full modification. It appears that at least 2 monomers, according to claim 1, are preferable to obtain the advantageous effect.

30

The modified PNA molecule according to the present invention comprises a PNA oligomer of a sequence, which is complementary to at least one target nucleotide sequence in the target cell. The target may be a nucleotide sequence of any RNA, which is essential for the growth, and/or reproduction of the cell.

35

The binding of a PNA strand to a DNA or RNA strand can occur in one of two orientations,


anti-parallel or parallel. As used in the present invention, the term complementary as applied to PNA does not in itself specify the orientation parallel or anti-parallel. It is significant that the most stable orientation of PNA/DNA and PNA/RNA is anti-parallel. In a preferred embodiment, PNA targeted to single strand RNA is complementary in an anti-parallel orientation.

The ability of PNAs to act as an antisense drug may be measured in many ways, which should be clear to the skilled person. To illustrate one way of preparing glycosylated PNA monomers with subsequent measuring of cell specificity, the following procedure may be used. However, the present invention is not limited hereto.

### Preparation of O-glycosylated PNAs

The initially employed strategy is outlined in Scheme 1. As orthogonal protection group for the synthesis of O-glycosylated PNAs, Fmoc, *tert*-butyl and allyl were applied for amine, hydroxyl and carboxylic acid protection, respectively.

Scheme 1




R=H) 1, 3,5,7,9,11,13,1s  
R=methyl) 2,4,6,8,10,12,14,2s

Initially, the carboxyl group of Fmoc-serine(t-Bu)-OH and Fmoc-threonine(t-Bu)-OH were allylated using allyl bromide in presence of DIPEA (18) to obtain high yields of **1-2**. Fmoc-deprotection by piperidinolysis treatment gave the free amine of **3-4** (19). The PNA backbone **5-6** were prepared using 2-Boc-aminoacetaldehyde (20) by reductive amination in presence of sodium cyanoborohydride. Subsequently, thymine-1-yl acetic acid (21) was condensed to the PNA backbone to give **7-8**. Boc and *tert*-butyl group removal by TFA resulted in the intermediates **9-10**. The ethereal solution was neutralised by addition of excess solid sodium carbonate. The free amine was re-protected with Fmoc using Fmoc-O-succinimide to obtain **11-12**.

Galactose donors were prepared by the method shown in scheme 2. Commercially available  $\beta$ -D-galactose pentaacetate was converted stereoselectively to the 1-O-deacetylated form **M1** by treatment with ammonia in a THF-methanol solution. The reaction was monitored by TLC, in order to avoid undesired further deacetylations, and **M1** was obtained in quantitative yield in the  $\alpha$ -form (16). Galactosyl trichloroacetimidate derivatives **M2** & **M3** were prepared by the method of Schmidt et. al. (17). *In situ* deprotonation of **M1** under basic condition ( $K_2CO_3$ ), reaction with trichloroacetonitrile followed by separation on silica gel column, gave  $\alpha$  and  $\beta$  anomers in good yields.

Scheme 2




A) ammonia in THF-methanol (7:3 v/v), 1.5 h, rt, ( $\approx 100\%$ ).

B)  $K_2CO_3$ ,  $CCl_3CN$ , mol. sieves.  $4A^\circ$ , DCM, over night, rt, purification by silicagel chromatography (hexane-EtOAc 2:1 v/v)  $\rightarrow \alpha$ : 36%,  $\beta$ : 48%

The reaction of both  $\alpha$ - and  $\beta$ -galactosyl trichloroacetimidates with **11** gave **13** as a 1:1 anomeric mixture, and reaction with **12** gave **14** predominantly as the  $\beta$ -form. The  $\beta$ -glycosidic linkage in **13** and **14** were verified by the  $^1H$  NMR triplets at  $\delta$  5.05 ppm and 5.08 ppm respectively ( $J_{1,2} = 7.7$  Hz). Finally, removal of the allyl group was performed in a high yield by treatment with *N*-ethyl aniline in the presence of catalytic amount of tetrakis ( $Ph_3P$ ) $_4$ Pd to give **1s-2s**.

## Preparation of lysine based N-C-glycosylated PNAs

Scheme 3




5

*N*-Fmoc-lysine-*N'*-(Dde)-OH was allylated using allyl bromide in presence of DIPEA and heating (18) resulting in **15**. Following Fmoc deprotection with 20% piperidine in DCM, the free amine of **16** was achieved. It has been reported that the protection group Dde is able to migrate from the side chain of lysine to an unprotected amine group of another lysine residue (22). Therefore, fresh **16** was used for the synthesis of the PNA backbone **17**. Subsequently, thymine-1-yl acetic acid (21) was condensed to the PNA backbone to yield **18**. The Dde protection group was removed by 2% hydrazine in DMF to give **19**. Allyl alcohol was added the deprotection solution to prevent reduction of the allyl group by hydrazine (23).

15

Scheme 4



5 A) Allyltrimethylsilane


B) OsO<sub>4</sub>, KIO<sub>4</sub>, dioxane-water (8:2 v/v), 12h, rt, (80%).

C) 5% aqueous NaH<sub>2</sub>PO<sub>4</sub>, 1M aqueous KMnO<sub>4</sub>, t-BuOH, 15 min, rt, (85%).

The sugar derivatives **M6** were prepared by standard literature methods (Scheme 4). Per-  
 10 acetylated galactose, mannose and fucose were converted to the corresponding α-C-allyl-  
 derivatives **M4** (24) and only α-epimers were isolated. **M4** were then oxidated to (α-D-)-  
 acetaldehydes **M5** (25) using potassium periodate in the presence of OsO<sub>4</sub> as catalyst. Fur-  
 further oxidation of the galactose and mannose derivatives by KMnO<sub>4</sub> gave the corresponding  
 acids **M6** (26). Attachment of 2 eq of **M5a** or **M5c** (galactose, fucose) derivatives to the free  
 15 amino group of **19**, and subsequent reduction by NaCNBH<sub>3</sub> gave good yields of **20-21**. The  
 reaction of **M6a-c** with **19** in the presence of DCC and DhbtOH gave **22-24**. These products  
 were converted to the Fmoc protected derivatives in two steps. They were first treated with  
 5% TES in TFA in order to remove the Boc group. The ethereal solution of the intermediates  
 were neutralised with excess solid sodium carbonate, and the free amine re-protected with  
 20 Fmoc using Fmoc-O-succinimide to obtain **25-27**. Removal of the allyl protection group from  
**20-22** and **25-27** yielded **3s-7s**.


## Preparation of N-glycosylated PNAs

Scheme 5



- 5 *N*-Fmoc-L-asp(OtBu)-OH was allylated to give **28**. The fmoc group was removed to give the free amine **29**. **29** reacted with fmoc-aminoacetaldehyde (**27**) to give **30**. Fmoc-aminoacetaldehyde was prepared by oxidation of fmoc-amino-2,3-propandiol (**28**) with potassium-*m*-periodate in dioxane-water (8:2 v/v). Thymine-1-ylacetic acid was attached to the backbone **30** to give **31**. *t*Butyl group on the side chain carboxylic acid was removed with TBAH in TES in TFA to give **32** in a moderate yield. All glycosylamines (scheme 6) were prepared using standard literature methods (**29**) via glycosyl-azides (**30**) and were obtained as  $\beta$ -epimers.
- 10

Scheme 6




**M7a)** R=Br, R'=OAc, galactose  
**M7b)** R=Br, R'=OAc, mannose  
**M7c)** R=Br, R'=OAc, fucose  
**M7d)** R=Cl, R'=NHAc, glucosamine  
**M7e)** R=Cl, R'=NHAc, galactosamine

Attachment of these sugar-amine compounds to the free carboxyl group of **32** was accomplished by the reaction of 2 eq of amine in presence of DCC and DhbtOH to give **33-37**. Final removal of the allyl groups was performed in a high yield whereby **9s-13s** were obtained.

### Preparation of C-galactosylated PNA

10 Scheme 7



The C-galactosylated PNA monomer was prepared from phosphonate **38** (Scheme 7). Alkaline hydrolysis of **38** in methanol afforded carboxylic acid **39**, which was subsequently converted to the t-Butyl ester **40** in 90% yield by treatment with EEDQ in t-butanol and chloroform for 24 h. Condensation of galactosyl-aldehyde derivative **M5** with phosphorylglycine t-butyl



ester **40** in the presence of *N,N,N',N'*-tetramethylguanidine as base gave **41** a mixture of E/Z isomers (5:95). As already pointed out by *U. Schmidt* and co-workers (31), the use of a strong base such as DBU or TMG favours the formation of the Z form. The E/Z ratio of **41** was measured to approximately 1:20 by proton NMR spectroscopy. Hydrogenation of **41** reduced the double bond and removed Cbz simultaneously, whereby intermediate **42** was obtained. This intermediate was subsequently treated with Fmoc-aminoacetaldehyde yielding **43** as a separable mixture of stereoisomers (25:75 according to TLC). The major diastereomer was readily isolated by flash chromatography. Attachment of thymine-1-yl acetic acid to the modified PNA backbone afforded **44**. Finally, the t-butyl group was removed by TFA scavenger and **14s** was obtained in a moderate yield.

### Solid Phase Synthesis

#### Procedure a) O-Glycosylated PNA

A well-characterised decamer PNA was applied as an antibacterial agent (32). The thymine-based PNA monomers of the decamer were replaced with O-galactosylated PNA monomers (**1s-2s**). The high lability of O-glycosidic bonds necessitated the use of a mild method for solid phase synthesis of O-glycosylated PNA oligomers. The Fmoc strategy was applied, since it has been adapted to O-glycosyl peptide solid phase synthesis (33). The stability of the o-galactosylated PNA to acidic cleavage conditions was tested on several resins. The Fmoc-Sieber-TG resin was found suitable due to its susceptibility to mild cleavage conditions (2% TFA in DCM). The  $\beta$ -elimination of the sugar moiety during synthesis and deprotection steps was also investigated. The HATU-collidine was found the most suitable coupling reagent. HATU was neutralised with collidine prior to acid pre-activation. The deprotection was carried out by anhydrous 50% morpholine in DMF at the minimum time required for deprotection (10+5 min). Coupling completion was investigated by Kaiser test and capping was avoided. As a cleavage mixture, 5% water, 30% TFA in DCM was sufficient to cleave the BHOC groups. Finally, a methanolic hydrazine solution (pH 9.5) was applied for the post-cleavage de-acetylation of the sugar-residue.

#### Procedure b) Lysine-based -C, N, C-glycosylated PNA

Initially, the Boc-strategy for the solid phase synthesis of PNA oligomers containing lysine based c-glycosylated-PNA monomers was applied. In the case of oligomers containing one residue of **s5**, following cleavage with TFMSA, a side product consisting of oligomer minus

one acetyl-group was detected by MALDI, and HPLC. Although the final deacetylation of both the product and the side product gave the target oligomer, it was decided to switch to the fmoc strategy, in order to avoid the use of strong acidic cleavage condition in boc strategy. For this purpose **22-24** was converted to the corresponding fmoc protected derivatives **5s-8s**. C- and N-glycosylated PNA monomers **9s-14s** were also prepared as fmoc protected derivatives. C- and N-glycosylated PNA monomers was also prepared as fmoc protected derivatives. PNA oligomers were synthesized on Fmoc-PAL-PEG resin. HATU-DIPEA was used as the coupling reagents and 20% piperidine in DMF at minimum required time for deprotections (3+2 min). Coupling yields were detected by Kaiser test and capping steps were avoided. A cleavage reagent of 5% TES in TFA was used and acetyl groups were removed with methanolic hydrazine (pH 10) solution following cleavage.

Several oligomers were synthesised and melting point temperature (T<sub>m</sub>) values were determined by standard methods. All of glycosylated PNAs showed reasonable binding affinity.

#### 15 **c) Glycosylated PNAs having cysteine at the N-terminal**

In order to develop the chemically conjugates of glycosylated PNA with biologically interesting compounds such as peptides or labelling compounds, it was necessary to optimise a method for the solid phase synthesis of the conjugate. For this purpose, cysteine was introduced at the N-terminal end of the glycosylated PNA as a S-trityl-protected derivative. Deprotection of sugar hydroxyl groups were performed on the resin prior to cleavage.

20 Eight decamers were prepared with incorporated glycosylated PNA monomers. The monomers were introduced in two or three residues shown as T\* in Table 1.

Table 1

**Cys-C-T\*\*\*-C-A-T\*\*-A-C-T\*-C-T-NH<sub>2</sub>**

PNA No:	Incorporated monomer	Number of residues
PNA: 1	7s	T*, T**
PNA: 2	8s	T*, T**
PNA: 3	9s	T*, T**
PNA: 4	10s	T*, T**
PNA: 5	11s	T*, T**
PNA: 6	12s	T*, T**, T***
PNA: 7	13s	T*, T**, T***
PNA: 8	14s	T*, T**, T***

5


### In-vivo imaging of glycosylated PNAs

Positron Emission Tomography (PET), a high-resolution sensitive and non-invasive imaging technique for the labelling of oligonucleotides (34) containing a single phosphorotioate monoester with an electrophilic moiety such as 2-bromo-N-substituted acetamides was applied. N-(4-halobenzyl)-2-bromoacetamide was designed as a radiochemically feasible reagent, the benzyl function offering the opportunity to act as the carrier of a radioactive halogen such as fluorine-18, the most widely used positron emitter ( $T_{1/2} = 109,8$  min.). Cysteine has a high and selective reactivity towards N-(4-fluorobenzyl)-2-bromoacetate, due to its nucleophilic thiol function. Based hereupon, eight glycosylated PNA decamers (shown in table 1) were prepared.

#### a) Synthesis of non-radioactive references

PNA monomers were conjugated with N-(4-fluorobenzyl)-2-bromoacetamide in a mixture of acetonitrile and phosphate buffer (0,1 M aq., pH 8.75) for 20 minutes at 60°C (scheme 8). The conjugated PNAs were purified by semi-preparative reverse phase HPLC and characterised by mass spectroscopy analysis (MALDI-TOF).

Scheme 8




5

## b) Radiochemistry

N-(4-[ $^{18}\text{F}$ ]fluorobenzyl)-2-bromoacetamide was synthesised in three steps using a robot. Typically, 60-90 mCi of pure N-(4-[ $^{18}\text{F}$ ]fluorobenzyl)-2-bromoacetamide was obtained in 85-95 min. starting from a 550-650 mCi of a cyclotron [ $^{18}\text{F}$ ]F $^-$  production batch (scheme 9).

10

Scheme 9



- 15 The HPLC-collected fraction containing N-(4-[ $^{18}\text{F}$ ]fluorobenzyl)-2-bromoacetamide was concentrated to dryness at 80°C under a nitrogen stream, before diluted with 0.4 ml acetonitrile. A solution of 1.3 mg PNA in 0.5 ml phosphate buffer (0.1 M, pH 8.75) and 0.1 ml of acetonitrile was rapidly added. The reactor was placed in a heating block and heated without stirring under a slight flow of nitrogen at 60°C for 20 minutes. Before total dryness, 1 ml distilled water was added to the reaction mixture, and the suspension was subjected to the HPLC purification for separation of labelled PNA[ $^{18}\text{F}$ ], unreacted PNA and unreacted N-(4-[ $^{18}\text{F}$ ]fluorobenzyl)-2-bromoacetamide. Labelled PNAs co-eluted with authentic synthesized unlabelled reference compounds. The HPLC fraction containing the labelled PNA was concentrated and formulated by transferring into a volume of 1-2 ml of serum.

### c) In-vivo PET imaging


Two male and two female Spargue-Dawley rats (200 g) were injected with 40 mCi labelled PNA in the tail vein placed in a Siemens ECAT EXACT HR+ camera under anesthesia and whole body images acquired in 3D mode for 2 hours.

### d) Ex-vivo study of bio-distribution

Following imaging, the animals were sacrificed and the kidneys, liver, spleen, heart, lungs, brain, muscles, blood and adrenals immediately collected. Aliquots of these organs were weighed and radioactivity was counted on a radiocounter. Radioactivity was expressed as percentage of injected dose per gram of organ (%ID/g) and reported as the mean  $\pm$  standard deviation.


Example of PNA oligomers containing glycosylated monomers used for pharmacokinetic analyses by  $^{18}\text{F}$ -isotope PET scanning is shown in Figure 1:

Figure 1



Organ distribution of PNA oligomers in rats is shown in Figure 2. The N-acetyl-galactosamine PNA shown in Figure 2 is preferentially targeting the liver. Similar results were obtained in baboons.


**5 Figure 2**



20

Pharmacokinetics of PNA oligomers of the type shown in figure 1 analysed by PET scanning. The N-acetyl-galactosamine PNA shown in Fig. 3 is preferentially accumulated in the liver. Similar results were obtained baboons.

**25      Figure 3**



e) Study of metabolism

- 5 Plasma supernatants were centrifuged in microfilter/10000 fixed Eppendorf tubes and subjected to analytical RP-HPLC. Urine samples were subjected to the HPLC directly. In some cases the samples were co-injected with unlabelled PNA.

10 **Pharmaceutical Compositions**

The PNA drugs of the present invention are used in the manufacture of medicaments for the treatment or prevention of bacterial, viral, protozoal, and fungal infections, cancer, metabolic diseases, cardiovascular diseases, autoimmune and immunological disorders, or for disinfecting non-living objects, such as surgery tools, hospital inventory, dental tools, slaughterhouse inventory and tool, dairy inventory and tools, barbers and beauticians tools and the like.

Within the present invention, the compounds of the invention may be prepared in the form of pharmaceutically acceptable salts, especially acid-addition salts, including salts of organic acid, fumaric acid, acetic acid, propionic acid, glycolic acid, lactic acid, pyruvic acid, oxalic acid, succinic acid, malic acid, tartaric acid, citric acid, benzoic acid, salicylic acid, and the like. Suitable inorganic acid-addition salts include salts of hydrochloric, hydrobromic, sulphuric- and phosphoric acids and the like. Further examples of pharmaceutically acceptable inorganic or organic acid addition salts include the pharmaceutically acceptable salts listed in Journal of Pharmaceutical Science, Berge et al 1977 (19), which are known to the skilled artisan.

Also intended as pharmaceutically acceptable acid-addition salts are the hydrates, which the present compounds are able to form.

The acid-addition salts may be obtained as the direct drugs of compound synthesis. In the alternative, the free base may be dissolved in a suitable solvent containing the appropriate acid, and the salt isolated by evaporating the solvent or otherwise separating the salt and solvent.

The compounds of this invention may form solvates with standard low molecular weight solvents using methods known to the skilled artisan.

In one aspect, the invention concerns the manufacture of a composition for treating or preventing bacterial, viral, protozoal, and fungal infections, cancer, metabolic diseases, cardiovascular diseases, autoimmune and immunological disorders, or disinfecting non-living objects, such as surgery tools, hospital inventory, dental tools, slaughterhouse inventory and tool, dairy inventory and tools, barbers and beauticians tools and the like.

Typical compositions include a compound of the invention or a pharmaceutically acceptable acid-addition salt thereof, associated with a pharmaceutically acceptable excipient which may be a carrier or a diluent or be diluted by a carrier, or enclosed within a carrier, which can be in the form of a capsule, sachet, paper or other container. In making the compositions, conventional techniques for the preparation of pharmaceutical compositions may be used. For example, the active compound will usually be mixed with a carrier, or diluted by a carrier, or enclosed within a carrier, which may be in the form of an ampoule, capsule, sachet, paper, or other container. When the carrier serves as a diluent, it may be solid, semi-solid, or liquid material, which acts as a vehicle, excipient, or medium for the active compound. The active compound can be adsorbed on a granular solid container for example in a sachet. Some examples of suitable carriers are water, salt solutions, alcohol's, polyethylene glycol's, polyhydroxyethoxylated castor oil, peanut oil, olive oil, glycine, gelatin, lactose, terra alba, sucrose, glucose, cyclodextrine, amylose, magnesium stearate, talc, gelatin, agar, pectin, acacia, stearic acid or lower alkyl ethers of cellulose, silicic acid, fatty acids, fatty acid amines, fatty acid monoglycerides and diglycerides, pentaerythritol fatty acid esters, polyoxyethylene, hydroxymethylcellulose and polyvinylpyrrolidone. Similarly, the carrier or diluent may include any sustained release material known in the art, such as glyceryl monostearate or glyceryl distearate, alone or mixed with a wax. The formulations may also include wetting agents, emulsifying and suspending agents, preserving agents, sweetening agents, thickeners or flavoring agents. The formulations of the invention may be formulated so as to provide quick, sustained, or delayed release of the active ingredient after administration to the patient by employing procedures well known in the art.

The pharmaceutical compositions can be sterilized and mixed, if desired, with auxiliary agents, emulsifiers, salt for influencing osmotic pressure, buffers and/or coloring substances and the like, which do not deleteriously react with the active compounds.

For therapeutic or prophylactic treatment, the PNA drug of the invention can be formulated in



a pharmaceutical composition, which may include one or more active ingredients such as antimicrobial agents, anti-inflammatory agents, anaesthetics, and the like in addition to PNA.

The pharmaceutical composition may be administered in a number of ways depending on whether local or systemic treatment is desired, and on the area to be treated. Administration may be done topically (including ophthalmically, vaginally, rectally, intranasally), orally, by inhalation, or parenterally, for example by intravenous drip or subcutaneous, intraperitoneal or intramuscular injection.

Formulations for topical administration may include ointments, lotions, creams, gels, drops, suppositories, sprays, liquids and powders. Conventional pharmaceutical carriers, aqueous, powder or oily bases, thickeners and the like may be necessary or desirable. Coated condoms may also be useful.

Compositions for oral administration include powders or granules, suspensions or solutions in water or non-aqueous media, capsules, sachets, or tablets. Thickeners, flavourings, diluents, emulsifiers, dispersing aids or binders may be desirable.

If a solid carrier is used for oral administration, the preparation may be tableted placed in a hard gelatin capsule in powder or pellet form or it can be in the form of a troche or lozenge. If a liquid carrier is used, the preparation may be in the form of a suspension or solution in water or a non-aqueous media, a syrup, emulsion or soft gelatin capsules. Thickeners, flavorings, diluents, emulsifiers, dispersing aids or binders may be added.

Formulations for parenteral administration may include sterile aqueous solutions, which may also contain buffers, diluents and other suitable additives.

For nasal administration, the preparation may contain a compound of the invention dissolved or suspended in a liquid carrier, in particular an aqueous carrier, for aerosol application. The carrier may contain additives such as solubilising agents, e.g. propylene glycol, surfactants, absorption enhancers such as lecithin (phosphatidylcholine) or cyclodextrine, or preservatives such as parabenes.

For parenteral application, particularly suitable are injectable solutions or suspensions, preferably aqueous solutions with the active compound dissolved in polyhydroxylated castor oil.

Tablets, dragees, or capsules having talc and/or a carbohydrate carrier or binder or the like are particularly suitable for oral application. Preferable carriers for tablets, dragees, or capsules include lactose, cornstarch, and/or potato starch. A syrup or elixir can be used in cases where a sweetened vehicle can be employed.

5

In yet another aspect, the invention concerns the treatment or prevention of bacterial, viral, protozoal, and fungal infections, cancer, metabolic diseases, cardiovascular diseases, autoimmune and immunological disorders, or treatment of non-living objects.

10 Dosing is dependent on severity and responsiveness of the condition to be treated, but will normally be one or more doses per day, with course of treatment lasting from several days to several months or until a cure is effected or a diminution of disease state is achieved. Persons of ordinary skill can easily determine optimum dosages, dosing methodologies and repetition rates.

15

Usually, dosage forms suitable for oral, nasal, pulmonal or transdermal administration comprise from about 0.01 mg to about 500 mg, preferably from about 0.01 mg to about 100 mg of the compounds of the invention admixed with a pharmaceutically acceptable carrier or diluent.

20 Treatments of this type can be practiced on a variety of organisms ranging from unicellular prokaryotic and eukaryotic organisms to multicellular eukaryotic organisms. Any organism that utilises DNA-RNA transcription or RNA-protein translation as a fundamental part of its hereditary, metabolic or cellular control is susceptible to therapeutic and/or prophylactic treatment in accordance with the invention. Seemingly diverse organisms such as bacteria,  
25 yeast, protozoa, algae, all plants and all higher animal forms, including warm-blooded animals, can be treated. Further, since each cell of multicellular eukaryotes can be treated since they include both DNA-RNA transcription and RNA-protein translation as integral parts of their cellular activity. Furthermore, many of the organelles, (e.g. mitochondria and chloroplasts) of eukaryotic cells also include transcription and translation mechanisms. Thus, single  
30 cells, cellular populations or organelles can also be included within the definition of organisms that can be treated with therapeutic or diagnostic PNA drug. As used herein, therapeutics is meant to include the eradication of a disease state, by killing an organism or by control of erratic or harmful cellular growth or expression.

35

## EXPERIMENTAL

The following examples are merely illustrative of the present invention and should not be considered limiting of the scope of the invention in any way.

5

### Abbreviations

The following abbreviations related to reagents are used in the experimental part:

aeg-PNA (Figure 4)	Amino Ethyl Glycine-PNA alias unmodified PNA
A monomer	N-(2-Boc-aminoethyl)-N-(N <sup>6</sup> -(benzyloxycarbonyl)adenine-9-yl-acetyl)glycine
Boc	Tert butyloxycarbonyl
Boc-Lys(2-Cl-Z)-OH	N- $\alpha$ -Boc-N- $\epsilon$ -2-chlorobenzyloxycarbonyl-L-lysine
C monomer	N-(2-Boc-aminoethyl)-N-(N <sup>4</sup> -(benzyloxycarbonyl)cytosine-1-yl-acetyl)glycine
TDBTU	2-(3,4-dihydro-4-oxo-1,2,3-benzotriazin-3-yl)-1,1,3,3-tetramethyluronium tetrafluoroborate
DIC	Diisopropylcarbodiimide
HOBt	N-Hydroxybenzotriazole
DCC	Dicyclohexylcarbodiimide
DCM	Dichloromethane
DIEA	N,N-diisopropylethylamine
DMF	N,N-dimethylformamide
DMSO	Dimethyl sulfoxide
G monomer	N-(2-Boc-aminoethyl)-N-(N <sup>2</sup> -(benzyloxycarbonyl)guanine-9-yl-acetyl)glycine
HATU	N-[(1-H-benzotriazole-1-yl)(dimethylamine)methylene]-N-methylmethanaminiumhexafluorophosphate N-oxide
HBTU	2-(1-H-benzotriazole-1-yl)-1,1,3,3-tetramethyluronium hexafluorophosphate
J monomer /nucleobase	N-(2-Boc-aminoethyl)-N-(N-2-(benzyloxycarbonyl) isocytosine-5-yl-acetyl)glycine
MBHA resin	p-methylbenzhydrylamine resin

NMP	N-methyl pyrrolidone
PTSA	Para - Toluene sulphonic Acid
T monomer	N-(2-Boc-aminoethyl)-N-(thymine-1-yl-acetyl)glycine
THF	Tetrahydrofuran
TFA	Trifluoroacetic acid
TFMSA	Trifluoromethanesulphonic acid
Tris	2-amino-2-(hydroxymethyl)-1,3-propanediol
Bn	Benzyl
Bz	Benzoyl
Ts	Tosyl
DMAP	4-dimethylaminopyridine
Tr	Trityl
DMEDA	Dimethylethylenediamine
EDC	N-(3-dimethylaminopropyl)-N'-ethylcarbodiimide
HOBt	1-Hydroxybenzotriazole
DIPEA	Diisopropylethylamine

Amino Acid	Abbr. I	Abbr. II	Side Chain Group
Alanine	A	Ala	methyl
Arginine	R	Arg	3-guanidinopropyl
Aspartic acid	D	Asp	carboxymethyl
Asparagine	N	Asn	aminocarboxymethyl
Cysteine	C	Cys	mercaptomethyl
Glutamic acid	E	Glu	2-carboxyethyl
Glutamine	Q	Gln	aminocarboxyethyl
Histidine	H	His	imidazol-4-yl-methyl
Isoleucine	I	Ile	1-methylpropyl
Leucine	L	Leu	2-methylpropyl
Lysine	K	Lys	4-aminobutyl
Methionine	M	Met	2-(methylthio)ethyl
Phenylalanine	F	Phe	benzyl
Serine	S	Ser	hydroxymethyl
Threonine	T	Thr	1-hydroxyethyl
Tryptophan	W	Try	3-indolyl

Amino Acid	Abbr. I	Abbr. II	Side Chain Group
Tyrosine	Y	Tyr	4-hydroxybenzyl
Valine	V	Val	1-methylethyl
Homoserine	-	(Hse)	2-hydroxymethyl
Citrulline	-	(Cit)	3-ureidopropyl
4-pyridyl-alanine	-	(4-Py)	4-pyridomethyl

Reagents and solvents were obtained from commercial sources and used without further purification, unless indicated. NMR spectra were recorded in CDCl<sub>3</sub> and CD<sub>3</sub>OH on Varian 400, 300 MHz and Bruker 250 MHz unity spectrometers, FAB mass spectra on a JEOL HX 110/110 mass spectrometer, MALDI-TOF mass spectra on a Cratos Compact Maldi II spectrometer. The Microanalysis Department of HCØ Institute, University of Copenhagen, performed elementary analyses. Flash chromatography was carried out using Silica Gel 60 (Merck particle size 0.040-0.063mm). The oligomers were analysed on a Delta Pak C<sub>18</sub> column (5µM, 3.9x150mm) and were purified on a Delta Pak C<sub>18</sub> column (15µM, 19x300mm). A gradient composed of A (0.1%TFA in water) and B (0.1% TFA in 10% H<sub>2</sub>O/90% acetonitrile) was used for analytical and preparative HPLC. Analytical: Time 0, 0% B. Time 35 min, 50% B (Flow, 1ml/min). Preparative: Time 0, 15% B. Time 45 min, 40% B (Flow, 8ml/min).

#### EXAMPLE 1

##### Procedure (a): allylation of N-protected amino acids

5 mmol of *N*-protected aminoacid was dissolved in a mixture of 10 ml acetonitrile and 12 ml allylbromide. (2.1 eq) *N*-*N*-diisopropylethylamine was added and the reaction mixture was stirred for 4 h at 40°C. Subsequently 200 ml ethylacetate was added and the solution was extracted with half saturated KHSO<sub>4</sub> (2 x 50 ml) and half saturated NaHCO<sub>3</sub> (2 x 50 ml) and 100 ml brine respectively. Organic phase was dried over MgSO<sub>4</sub> and volatile were removed under vacuum. The remaining oil was used for the next step (Yield = 87%-95%).

#### EXAMPLE 2

##### Procedure (b): preparation of N-boc/fmoc protected PNA backbone

12 mmol amine and 10 mmol boc/fmoc-aminoacetaldehyde were stirred in 40 ml methanol for 10 min. 0.8 ml (13 mmol) Acetic acid and 0.6 g (10 mmol) sodium cyanoborohydrate was added sequentially. The reaction mixture was stirred for further 1 h at room temperature. Volatiles was removed under vacuum and the remaining solid dissolved in 100 ml ethylacetate and extracted with NaHCO<sub>3</sub> saturated solution and brine. Organic phase was dried over

magnesium sulphate and evaporated to dryness under vacuum. The residue was purified on silica gel column eluting hexane-ethylacetate 1:1. (Yield = 50% - 80%).

### EXAMPLE 3

#### 5 **Procedure (c): coupling of thymine-1-yl-acetic acid to the PNA backbone**

17 mmol Thymine-1-ylacetic acid and 18 mmol DhbtOH were dissolved in 50 ml dry DMF. 20 mmol DCC was added and the solution stirred for 20 minutes. 8.5 mmol backbone was added in 20 ml dry DMF to the reaction mixture and stirred for further 6 h at room temperature. Volatiles were removed under vacuum and the remaining dissolved in 200 ml ethylacetate. Insoluble DCU was filtered off and the filtrate extracted with NaHCO<sub>3</sub> saturated solution (2 x 100 ml) and brine (100 ml). After drying over magnesium sulphate, the organic phase was evaporated to dryness. The residue was purified on a silica gel column eluting the mixture of ethylacetate-methanol 10:0 to 10:1 (Yield = 63%-87%).

### 15 EXAMPLE 4

#### **Procedure (d): boc de-protection and subsequent fmoc protection**

3 mmol boc-protected compound was added to 20 ml solution of 5% triethylsilane in trifluoroacetic acid at 0°C and stirred until TLC did not show any starting material left. 50 ml Toluene was added and volatile removed under vacuum. Further 3 x 100 ml DCM was added and evaporated in order to removal of TFA. The remaining was dissolved in 50 ml diethylether and 5 g sodium carbonate was added as well powdered solid. The suspension was stirred for 30 min and then evaporated. The solid residue was suspended in 50 ml Acetonitrile and 3.2 mmol fmoc-O-Su was added and stirred for further 2h. The solution was evaporated under vacuum and the crude was purified on silica gel column eluting ethylacetate-methanol 10:0 to 10:1 (Yield = 88%-94%).

### EXAMPLE 5

#### **Procedure (e): N-glycosylation**

2 mmol **32**, 2.2 mmol DhbtOH, and 3 mmol DCC were mixed in 10 ml DMF and stirred for 30 min under Nitrogen. A solution 2 mmol Sugar-amine in 10 ml DMF was added and the reaction mixture was stirred overnight. Volatile was removed under vacuum and the residue dissolved in 200 ml ethylacetate. Insoluble DCU was filtered off and the filtrate extracted with NaHCO<sub>3</sub> saturated solution (2 x 100ml) and brine (100 ml). After drying over magnesium sulphate the organic phase was evaporated to dryness. The residue subjected to silica gel column eluting the mixture of ethylacetate-methanol 10:0 to 10:1 (Yields = 60%-77%).

## EXAMPLE 6

**Procedure (f): removing of allyl group**

0.1 mmol of ester was dissolved in 2 ml THF. 23 mg (0.02 mmol, 0.2 eq)  
 5 Tetrakis(triphenylphosphine)Pd(0) was added. 10 eq N-ethyl-aniline was added drop wise to the reaction mixture and stirred at room temperature. The reaction was checked by TLC (ethylacetate-methanole 10:1). After complete conversion of starting material, the reaction mixture was poured dropwise in a 20 ml vigorously stirring solution of diethylether-n-hexane 1:1. White precipitate was collected by filtration and washed with n-hexane (Yield = 80%-  
 10 95%).

## EXAMPLE 7

**Threonine(t-Bu)-Allyl (4)**

To 4.4 g (10 mmol) **2** [prepared by *procedure (a)*] was added 50 ml of a solution of 20%  
 15 piperidine in DCM. After 30 min, 100 ml toluene was added and volatile were removed under vacuum. The residue was purified with a short silica gel column eluting ethylacetate-methanole 10:0 to 10:1. 1.6 g (7.5 mmol) titled compound was obtained as colourless oil (Yield = 75%).

[ $\alpha$ ] obtained as colourless oil in 83% yield.

20 [ $\alpha$ ]<sub>D</sub><sup>22</sup> = -1.53 (c = 1, methanol); MS (FAB) *m/z* 216 (M+H); <sup>1</sup>H-NMR, (CDCl<sub>3</sub>):  $\delta$  5.94 (9 line m, 1H, -CH=CH<sub>2</sub>), 5.3 (dd, 1H, *J*<sub>trans</sub> = 17.03, 1.37 Hz), 5.2 (dd, 1H, *J*<sub>cis</sub> = 10.5, 1.1 Hz), 4.64-4.46 (dddt, 2H, *J* = 33.08, 10.7, 5.8, 1.1, O-CH<sub>2</sub>-CH=), 4 (m, 1H, <sup>β</sup>CH), 3.26 (d, 1H, *J*=3.3 Hz, <sup>α</sup>CH), 1.65 (s, 2H, -NH<sub>2</sub>), 1.2 (d, 3H, *J*=6.3 Hz, <sup>β</sup>C-CH<sub>3</sub>), 1.09 (s, 9H, boc); <sup>13</sup>C-NMR (CDCl<sub>3</sub>):  $\delta$  174.51 (COO-Allyl), 132.09 (-CH=CH<sub>2</sub>), 118.87 (-CH=CH<sub>2</sub>), 73.77 [-O-C(Me)<sub>3</sub>], 68.63,  
 25 65.67, 60.84, 28.68 [-COO-C(CH<sub>3</sub>)<sub>3</sub>], 20.94 (<sup>β</sup>C-CH<sub>3</sub>).

C.H.N analyse for C<sub>11</sub>H<sub>21</sub>NO<sub>3</sub>. ½ H<sub>2</sub>O: calc. C 60.11, H 9.86, N 6.37; found C 59.87, H 9.95, N 6.34.

## EXAMPLE 8

**N-(2-Boc-aminoethyl)-Serine(t-Bu)-Allyl (5)**

*Preparation: Procedure (b)*

[ $\alpha$ ]<sub>D</sub><sup>22</sup> = -10.26 (c=1, methanol); MS (FAB) *m/z* 345(M+H); <sup>1</sup>H-NMR, (CDCl<sub>3</sub>):  $\delta$  5.96-5.89 (8  
 line m, 1H, -CH=CH<sub>2</sub>), 5.26 (dd, 1H, *J*<sub>trans</sub>=17.2, 1.46 Hz), 5.17 (dd, 1H, *J*<sub>cis</sub>=10.4, 1.28 Hz),  
 5.05 (br.s, boc-HN-), 4.57-4.55 (m, 2H, O-CH<sub>2</sub>-CH=), 3.52 (ABq, 2H, *J*=4.95Hz, <sup>β</sup>CH), 3.36 (t,  
 35 *J*=4.76 Hz, 1H, <sup>α</sup>CH), 3.2-3.1 (m, 2H), 2.8 (m, 1H), 2.6 (m, 1H), 1.37 (s, 9H, *t*-butyl), 1.08 (s,

9H, boc);  $^{13}\text{C}$ -NMR ( $\text{CDCl}_3$ ):  $\delta$  172.50 (COO-Allyl), 155.9 (-NH-COO), 131.74 ( $-\underline{\text{CH}}=\text{CH}_2$ ), 118.42 ( $-\text{CH}=\underline{\text{CH}}_2$ ), 118.27, 73.13 [ $-\text{O}-\underline{\text{C}}(\text{Me})_3$ ], 65.263 ( $-\underline{\text{CH}}_2-\text{CH}=\text{}$ ), 62.66 ( $\beta$  carbon), 61.22 ( $\alpha$  carbon), 47.18, 40.03, 28.25 [ $-\text{COO}-\text{C}(\underline{\text{CH}}_3)_3$ ], 27.17 [ $-\text{O}-\text{C}(\underline{\text{CH}}_3)_3$ ].

C.H.N analyse for  $\text{C}_{17}\text{H}_{32}\text{N}_2\text{O}_5 \cdot \frac{1}{2} \text{H}_2\text{O}$ : calc. C 57.77, H 9.41, N 7.93; found C 58.11, H 9.20, N 8.14.

#### EXAMPLE 9

##### ***N*-(2-Boc-aminoethyl)-Threonine(*t*-Bu)-Allyl (6)**

*Preparation: Procedure (b)*

(Yield = 67%)  $[\alpha]_D^{22} = -3.39$  ( $c=1$ , methanol); MS (FAB)  $m/z$  359 ( $M+H$ );  $^1\text{H}$ -NMR, ( $\text{CDCl}_3$ ):  $\delta$  5.90-5.83 (8 line m, 1H,  $-\underline{\text{CH}}=\text{CH}_2$ ), 5.26 (dt, 1H,  $J_{\text{trans}}=17$ , 1.4 Hz), 5.18 (dt, 1H,  $J_{\text{cis}}=10$ , 1.1 Hz), 5.05 (br.s, boc-HN-), 4.60-4.47 (dddt, 2H,  $J=33.3$ , 13.2, 5.8, 1.3 Hz,  $\text{O}-\underline{\text{CH}}_2-\text{CH}=\text{}$ ), 3.9 (q, 1H,  $^{\beta}\text{CH}$ ), 3.14-3.08 (dm, 2H), 3.05 (d, 1H,  $J=3.5$ ,  $^{\alpha}\text{CH}$ ), 2.8 (m, 1H), 2.7 (m, 1H), 1.37 (s, 9H, *t*-butyl), 1.18 (d, 3H,  $J=6.2$  Hz,  $^{\beta}\text{C}-\text{CH}_3$ ), 1.06 (s, 9H, boc);  $^{13}\text{C}$ -NMR ( $\text{CDCl}_3$ ):  $\delta$  172.91 (COO-Allyl), 155.88 (-NH-COO), 131.65 ( $-\underline{\text{CH}}=\text{CH}_2$ ), 118.54 ( $-\text{CH}=\underline{\text{CH}}_2$ ), 78.81 [ $-\text{COO}-\underline{\text{C}}(\text{Me})_3$ ], 73.66 [ $-\text{O}-\underline{\text{C}}(\text{Me})_3$ ], 68.14 ( $\beta$  carbon), 66.37 ( $\alpha$  carbon), 65.270 ( $-\underline{\text{CH}}_2-\text{CH}=\text{}$ ), 47.51, 39.88, 28.23 [ $-\text{COO}-\text{C}(\underline{\text{CH}}_3)_3$ ], 28.19 [ $-\text{O}-\text{C}(\underline{\text{CH}}_3)_3$ ], 20.49 ( $^{\beta}\text{C}-\underline{\text{CH}}_3$ ).

C.H.N analyse for  $\text{C}_{18}\text{H}_{34}\text{N}_2\text{O}_5$ : calc. C 60.31, H 9.56, N 7.81; found C 60.16, H 9.98, N 7.75.

#### 20 EXAMPLE 10

##### ***N*-(2-Boc-aminoethyl)-*N*-(Thymine-1-ylacetyl)Serine(*t*-Bu)-Allyl (7)**

*Preparation: Procedure (c)*

Mp = 70-72; MS (FAB)  $m/z$  216 ( $M+H$ );  $^1\text{H}$ -NMR, ( $\text{CDCl}_3$ ): (major rotamer)  $\delta$  9.08 (s, 1H, Thymine aromatic), 6.83 (s, 1H, Thymine-NH-), 5.87-5.77 (8 line m, 1H,  $-\underline{\text{CH}}=\text{CH}_2$ ), 5.6 (br.s, 1H, boc-HN-), 5.25 (dd, 1H,  $J_{\text{trans}}=17.2$ , 1.28 Hz), 5.17 (dd, 1H,  $J_{\text{cis}}=10.2$ , 0.91 Hz), 4.6-4.4 (dm, 2H,  $-\text{O}-\underline{\text{CH}}_2-\text{CH}=\text{}$ ), 4.55 (s, 2H,  $-\text{CO}-\text{CH}_2\text{-Thymine}$ ), 4.26 (br.d, 1H,  $J=5.3$ ), 3.93 (t, 1H), 3.77 (dd, 1H,  $J=10$ , 3.1 Hz), 3.6-3.4 (dm, 2H), 3.35 (m, 2H), 1.83 (s, 3H, Thymine- $\text{CH}_3$ ), 1.37 (s, 9H, *t*-butyl), 1.1 (s, 9H, boc);  $^{13}\text{C}$ -NMR ( $\text{CDCl}_3$ ):  $\delta$  168.93, 167.15, 164.07, 155.85 (-NH-COO), 150.72, 140.72, 131.36 ( $-\underline{\text{CH}}=\text{CH}_2$ ), 118.84 ( $-\text{CH}=\underline{\text{CH}}_2$ ), 111.59, 110.43, 79.62 [ $-\text{COO}-\underline{\text{C}}(\text{Me})_3$ ], 73.84 [ $-\text{O}-\underline{\text{C}}(\text{Me})_3$ ], 66.06 ( $-\underline{\text{CH}}_2-\text{CH}=\text{}$ ), 60.79, 60.24, 59.50, 38.84, 33.66, [28.30, 28.10,  $-\text{COO}-\text{C}(\underline{\text{CH}}_3)_3$ ], 27.15 [ $-\text{O}-\text{C}(\underline{\text{CH}}_3)_3$ ], 12.25 (Thymine- $\text{CH}_3$ ).

#### EXAMPLE 11

##### ***N*-(2-Boc-aminoethyl)-*N*-(Thymine-1-ylacetyl)Threonine(*t*-Bu)-Allyl (8)**

*Preparation: Procedure (c)*



Mp = 71-73; Yield = 76%; MS (FAB)  $m/z$  525(M+H);  $^1\text{H-NMR}$ , ( $\text{CDCl}_3$ ): (major rotamer)  $\delta$  8.59 (s, 1H, Thymine aromatic), 6.9 (s, 1H, Thymine-NH-), 5.95-5.85 (8 line m, 1H, -CH=CH<sub>2</sub>), 5.5 (br.s, 1H, boc-HN-), 5.3 (dd, 1H,  $J_{\text{trans}}=17.1, 1.4$  Hz), 5.25 (d, 1H,  $J_{\text{cis}}=10.2$  Hz), 4.66 (s, 2H, -CO-CH<sub>2</sub>-Thymine), 4.52-4.41 (overlapping m, 3H, -O-CH<sub>2</sub>-CH=,  $^u\text{CH}$ ), 4.1 (q, 1H), 3.8 (dt, 1H,  $J=15.3, 6$  Hz), 3.6 (m, 1H), 3.5-3.4 (m, 2H), 1.89 (s, 3H, Thymine-CH<sub>3</sub>), 1.44 (s, 9H, *t*-butyl), 1.28 (d, 3H,  $J=6\text{Hz}$ ,  $^{\beta}\text{C-CH}_3$ ), 1.1 (s, 9H, boc);  $^{13}\text{C-NMR}$  ( $\text{CDCl}_3$ ): (major rotamer)  $\delta$  169.12, 167.95, 163.78, 155.78 (-NH-COO), 150.59, 140.73, 131.20 (-CH=CH<sub>2</sub>), 119.41 (-CH=CH<sub>2</sub>), 110.43, 79.61 [-COO-C(Me)<sub>3</sub>], 74.55 [-O-C(Me)<sub>3</sub>], 66.14 (-CH<sub>2</sub>-CH=), 64.39, 39.16, 33.79, [28.69, 28.60, 28.40, -COO-C(CH<sub>3</sub>)<sub>3</sub>], [25.53, 24.87, -O-C(CH<sub>3</sub>)<sub>3</sub>], 21.71 ( $^{\beta}\text{C-CH}_3), 12.37 (Thymine-CH<sub>3</sub>).$

#### EXAMPLE 12

##### ***N*-Fmoc-aminoethyl-*N*-(Thymine-1-ylacetyl)Serine(OH)-Allyl (11)**

*Preparation: Procedure (d)*

Mp = 101-103;  $[\alpha]_{\text{D}}^{22} = -5.36$  ( $c=0.25$ , methanol); MS (FAB)  $m/z$  577(M+H);  $^1\text{H-NMR}$ , ( $\text{CDCl}_3$ ):  $\delta$  9.63 (s, 1H, Thymine aromatic), 7.56-7.07 (8H, Florenyl aromatic protons), 6.59 (s, 1H, Thymine-NH-), 5.9 (s, 1H, fmoc-HN-), 5.8-5.6 (m, 1H, -CH=CH<sub>2</sub>), 5.12 (d, 1H,  $J_{\text{trans}}=17.2$  Hz), 5.05 (dd, 1H,  $J_{\text{cis}}=10.4, 0.91$  Hz), 4.6-4.4 (m, 3H, O-CH<sub>2</sub>-CH=,  $^u\text{CH}$ ), 4.24 (s, 2H, -CO-CH<sub>2</sub>-Thymine), 4-3.8 (m, 4H), 3.4-3.2 (m, 4H), 2.73 (br.s, 1H), 1.7 (s, 3H, Thymine-CH<sub>3</sub>);  $^{13}\text{C-NMR}$  ( $\text{CDCl}_3$ ):  $\delta$  168.50, 168.06, 167.94, 167.07, 164.05, 162.58, 156.59, 151.33, 143.58 and 143.39 (florenyl), 140.98, 131.14 (-CH=CH<sub>2</sub>), [(130.69, 127.54, 126.88, 124.82, 124.36, 119.75 (florenyl)], 118.95 (-CH=CH<sub>2</sub>), 110.90, 66.75, 66.17, 63.43 (-CH<sub>2</sub>-CH=), 59.47, 49.24, 48.81, 47.10, 39.45, 31.54, 12.30 (Thymine-CH<sub>3</sub>).

C.H.N analyse for C<sub>30</sub>H<sub>32</sub>N<sub>4</sub>O<sub>8</sub> · ½ H<sub>2</sub>O: calc. C 61.35, H 5.68, N 9.57; found. C 61.39, H 5.53, N 9.36.

#### EXAMPLE 13

##### ***N*-Fmoc-*N*-(Thymine-1-ylacetyl)Threonine(OH)-Allyl (12)**

*Preparation: Procedure (d)*

Mp = 99-101; Yield = 91%;  $[\alpha]_{\text{D}}^{22} = -0.93$  ( $c=0.75$ , methanol); MS (FAB)  $m/z$  591 (M+H);  $^1\text{H-NMR}$ , ( $\text{CDCl}_3$ ):  $\delta$  8.9 (s, 1H, Thymine aromatic), 7.6-7.1 (8H, Florenyl aromatic protons), 6.6 (s, 1H, Thymine-NH-), 5.8-5.6 (overlapping m, 2H, -CH=CH<sub>2</sub>, fmoc-HN-) 5.12 (dt, 1H,  $J_{\text{trans}}=16.1, 8.8$  Hz), 5.07 (dd, 1H,  $J_{\text{cis}} = 10.3, 1$  Hz), 4.5 (overlapping m, 3H), 4.42-4.26 (m, 2H, O-CH<sub>2</sub>-CH=), 4.2 (t, 1H,  $J=7.4$  Hz) 4.03 (d, 2H,  $J=6$  Hz), 3.65 (br.d, 1H,  $J=5.3$  Hz), 3.53-3.47 (m, 1H), 3.4-3.24 (m, 3H), 2.8 (br.s, 1H), 1.68 (s, 3H, Thymine-CH<sub>3</sub>), 1.14 (d, 3H,  $J=5.86$

Hz,  $^1\text{C}-\text{CH}_3$ );  $^{13}\text{C}$ -NMR ( $\text{CDCl}_3$ ):  $\delta$  168.67, 167.85, 163.99, 156.60, 151.10, [143.67, 143.50, 141.15, 140.98 (floreanyl)], 131.22 ( $-\text{CH}=\text{CH}_2$ ), 127.64, 126.96, 124.88, 124.36, 119.87, 119.25 ( $-\text{CH}=\text{CH}_2$ ), 110.75, 67.50, 66.61, 66.19, 65.16, 49.55, 48.86, 47.06, 39.41, 21.30 ( $^1\text{C}-\text{CH}_3$ ), 12.30 (Thymine- $\text{CH}_3$ ).

- 5 C.H.N analyse for  $\text{C}_{31}\text{H}_{34}\text{N}_4\text{O}_8 \cdot \frac{1}{2} \text{H}_2\text{O}$ : calc. C 62.09, H 5.88, N 9.34; found C 62.10, H 5.86, N 9.22.

#### EXAMPLE 14

##### ***N*-Fmoc-*N*-(Thymine-1-ylacetyl)Serine(2,3,4,5-tetra-*O*-acetyl- $\alpha$ -D-Galactose-1-yl)-Allyl (13)**

*Preparation:* 0.7g (1.2 mmol) **11** and 0.75 g (1.5 mmol) galactose trichloroacetamide were dissolved in 5 ml ethylacetate. 0.5 g molecular sieve was suspended and 1.2 ml (9.5 mmol) borontriflate in 1 ml ethylacetate was added at 0°C. Ice bath was removed and the reaction mixture was stirred under nitrogen over night. 50 ml ethylacetate was added and the reaction mixture was filtered over celite. The filtrate was extracted with an ice cold half saturated NaHCO<sub>3</sub> aqueous solution (2 x 25 ml), and brine (50 ml), dried over magnesium sulphate and evaporated to dryness. The residue was purified on silica gel column eluting hexane-ethylacetate 1:1. 0.4 g (0.9 mmol)  $\alpha$  anomer was obtained as white crystalline. (yield = 37%)

Mp = 115-118;  $[\alpha]_D^{22} = -24.17$  ( $c=0.5$ , methanol); MS (FAB)  $m/z$  907(M+H);  $^1\text{H}$ -NMR, ( $\text{CDCl}_3$ ):  $\delta$  8.75 (s, 1H, Thymine aromatic), 7.68-7.21 (m, 8H, Florenyl aromatic protons), 6.8 (s, 1H, Thymine-NH-), 5.9-5.7 (overlapping m, 2H,  $-\text{CH}=\text{CH}_2$ , fmoc-HN-), 5.3(d, 1H,  $J=2.9$  Hz, sugar  $^4\text{CH}$ ), 5.25 (d.d, 1H,  $J_{\text{trans}} = 17.2, 1.26$  Hz), 5.15 (d, 1H,  $J_{\text{cis}}=10.7$  Hz), 5.05 (t, 1H,  $J_{1,2}=7.7$  Hz,  $J_{2,3}=10.4$  Hz, sugar  $^2\text{CH}$ ), 4.95 (dd, 1H,  $J=10.4, 3.2$  Hz, sugar  $^3\text{CH}$ ), 4.7-4.3 (m, 6H), 4.25, 3.95 (m, 6H), 3.8 (t, 2H,  $J=6.4$ , sugar  $^5\text{CH}$ ), 3.6-3.3 (m, 3H), 2.9 (br.s, 1H), 2.1-1.9 (4s, 12H, sugar acetyl protons), 1.8 (s, 3H, Thymine- $\text{CH}_3$ );  $^{13}\text{C}$ -NMR ( $\text{CDCl}_3$ ):  $\delta$  169.37, 169.09, 168.96, 168.80, 166.84, 166.61, 163.09, 155.81(-NH-COO), 149.67, 142.76, 140.27, 130.37 ( $-\text{CH}=\text{CH}_2$ ), 126.73, 126.06, 123.91, 118.98, 118.24 ( $-\text{CH}=\text{CH}_2$ ), 109.26, 100.03 (sugar C1), 70.09, 69.57, 67.75, 66.38, 65.93, 65.54, 65.43, 60.22, 60.09, 47.48, 47.02, 46.29, 38.60, 19.78-19.54 (sugar-CO- $\text{CH}_3$  groups), 11.28 (Thymine- $\text{CH}_3$ ).

HRMS (M+Na)<sup>+</sup>, calculated (found) for  $\text{C}_{44}\text{H}_{50}\text{N}_4\text{O}_{17}\text{Na}$  are 929.3069 (929.3087).

#### EXAMPLE 15

##### ***N*-Fmoc-*N*-(Thymine-1-ylacetyl)Threonine(2,3,4,5-tetra-*O*-acetyl- $\alpha$ -D-Galactose-1-yl)-Allyl (14)**

*Preparation:* The Procedure is the same as described above for the synthesis of **13**

Yield = 77%, mp = 106-108;  $[\alpha]_D^{22} = -27.38$  ( $c=0.5$ , methanol); MS (FAB)  $m/z$  921(M+H);  $^1\text{H-NMR}$ , ( $\text{CDCl}_3$ ):  $\delta$  8.54 (s, 1H, Thymine aromatic), 7.78-7.76 (8H, Florenyl aromatic protons), 6.95 (1H, Thymine-NH-), 5.92-5.82 (m, 1H,  $-\text{CH}=\text{CH}_2$ ), 5.81 (s, 1H, fmoc-HN-), 5.38 (d, 1H,  $J=2.5$  Hz, sugar  $^4\text{CH}$ ), 5.3 (d, 1H,  $J_{\text{trans}}=17.4$  Hz), 5.2 (d, 1H,  $J_{\text{cis}}=10.4$  Hz), 5.08 (t, 1H,  $J_{1,2}=7.7$  Hz,  $J_{2,3}=10.4$  Hz, sugar  $^2\text{CH}$ ), 5.03 (dd, 1H,  $J=10.4$ , 3.3 Hz, sugar  $^3\text{CH}$ ), 4.9 (d, 1H,  $J=16.3$ ), 4.7-4.5 (overlapping s ddd and s, 4H,  $J=49.6$ , 13.2, 5.86 Hz,  $-\text{O}-\text{CH}_2-\text{CH}$  and  $\text{CO}-\text{CH}_2-\text{Thymine}$ ), 4.44 (overlapping, 2H), 4.35 (d, 2H,  $J=16.3$  Hz), 4.2 (t, 2H,  $J=7$  Hz), 4.07 (dd, 1H,  $J=22.9$ , 6.41 Hz, sugar  $^6\text{CH}$ ), 3.88 (dd, 1H,  $J_{5,6}=6.23$  Hz,  $J_{5,8}=6.59$ , sugar  $^5\text{CH}$ ), 3.6 (overlapping s, 2H), 3.43 (br.s, 1H), 2.18-2 (3s, 12H, sugar acetyl protons), 1.88 (s, 3H, Thymine- $\text{CH}_3$ ), 1.27 (d, 3H,  $J=5.86$ ,  $^8\text{C}-\text{CH}_3$ );  $^{13}\text{C-NMR}$  ( $\text{CDCl}_3$ ):  $\delta$  170.11, 169.89, 167.97, 163.65, 156.64(-NH-COO), 150.35, 143.53, 141.21, 141.05, 131.31( $-\text{CH}=\text{CH}_2$ ), 127.51, 126.82, 124.78, 119.80, 118.78 ( $-\text{CH}=\text{CH}_2$ ), 110.07, 98.73 (sugar C1), 73.34, 71.53, 70.70, 70.48, 70.01, 68.80, 66.66, 66.55, 66.17, 65.65, 64.10, 62.79, 60.84, 60.31, 48.48, 47.19, 39.70, [21.05, 20.83, 20.67, 20.55 (sugar-CO- $\text{CH}_3$  groups)], 17.38, 14.20 ( $^8\text{C}-\text{CH}_3$ ), 12.29 (Thymine- $\text{CH}_3$ ).

C.H.N analyse for  $\text{C}_{45}\text{H}_{52}\text{N}_4\text{O}_{17}$ .  $\text{H}_2\text{O}$ : calc. C 57.56, H 5.80, N 5.97; found C 57.51, H 5.52, N 5.82.

#### EXAMPLE 16

##### Lysine(Dde)-Allyl (16)

*Preparation:* 5.7g (10 mmol) ester **15** [prepared by *Procedure (a)*] was added to a solution of 20% piperidine in DCM and stirred for 30 min. 100 ml toluene was added and volatile were removed under vacuum. The residue was purified on a short silicagel column eluting ethylacetate-methanol 10:0 to 10:1. 2.5 g (7 mmol) Titled compound was obtained as slightly yellow oil which was used subsequently for the next step (Yield = 71%).

MS (FAB)  $m/z$  351(M+H);  $^1\text{H-NMR}$ , ( $\text{CDCl}_3$ ):  $\delta$  13.34(s, 1H, -HN-Dde), 5.9-5.8 (ddt, 1H,  $J=5.57$ , 10.55, 10.43 Hz,  $-\text{CH}=\text{CH}_2$ ), 5.27(dd, 1H,  $J_{\text{trans}}=17.3$ , 1.47 Hz), 5.22(dd, 1H,  $J_{\text{cis}}=10.5$ , 1.17 Hz), 4.56 (dt, 5.86, 1.17Hz,  $\text{O}-\text{CH}_2-\text{CH}=\text{}$ ), 3.4 (dd, 1H,  $J=7.5$ , 5 Hz,  $^{\alpha}\text{CH}$ ), 3.3 (ABq, 2H,  $J_{\text{AB}}=12.3$ ,  $-\text{CH}_2-\text{NH-Dde}$ ), 2.49 (s, 3H, -HN-C- $\text{CH}_3$ ), 2.3 (s, 4H, 2x- $\text{CH}_2-$  of Dde), 1.77 (s, 2H, - $\text{NH}_2$ ), 1.74-1.38 (overlapping m, 6H), 0.97 (s, 6H, 2x- $\text{CH}_3$  of Dde);

$^{13}\text{C-NMR}$  ( $\text{CDCl}_3$ ):  $\delta$  [198.59, 196.83 (-CO- of Dde)], 175.19 ( $=\text{C}(\text{CH}_3)-\text{NH}$ ), 173.14 (COO-Allyl), 131.60 ( $-\text{CH}=\text{CH}_2$ ), 118.51 ( $-\text{CH}=\text{CH}_2$ ), 107.56 ( $=\text{C}=\text{}$  of Dde), 65.33 ( $\text{O}-\text{CH}_2-\text{CH}=\text{}$ ), 60.10 ( $-\text{CH}_2-$  of Dde), 53.92, 42.96, 33.98, 29.83, 28.59, 28.02, 22.87, 17.65 ( $\text{CH}_3$  of Dde).

#### EXAMPLE 17

**N-(2-Boc-aminoethyl)-Lysine(Dde)-Allyl (17)***Preparation: Procedure (b), yield= %65*

MS (FAB)  $m/z$  494 (M+H);  $^1\text{H-NMR}$ , ( $\text{CDCl}_3$ ):  $\delta$  13.45 (s, 1H, -HN-Dde), 5.98-5.85 (10 line m, -CH=CH<sub>2</sub>), 5.33 (dd, 1H,  $J_{\text{trans}}=17.29$ , 1.47 Hz), 5.28 (dd, 1H,  $J_{\text{cis}}=10.55$ , 1.17 Hz), 4.56 (dd, 5.86, 1.17Hz, O-CH<sub>2</sub>-CH=), 3.4 (overlapping m, 3H,  $^{\circ}\text{CH}$ , -CH<sub>2</sub>-NH-Dde), 3.2 (m, 2H), 2.8 (m, 1H), 2.7 (m, 1H), 2.55 (s, 3H, -HN-C-CH<sub>3</sub>), 2.36 (s, 4H, 2x-CH<sub>2</sub>- of Dde), 1.8-1.5 (2 x m, 6H), 1.44 (s, 9H, boc), 1.03 (s, 6H, 2x-CH<sub>3</sub> of Dde);  $^{13}\text{C-NMR}$  ( $\text{CDCl}_3$ ):  $\delta$  [198.79, 197.21 (-CO- of Dde)], 174.68 (=C(CH<sub>3</sub>)-NH), 173.68 (COO-Allyl), 156.37 [-NH-COO-C(Me)<sub>3</sub>], 132.00 (-CH=CH<sub>2</sub>), 119.27(-CH=CH<sub>2</sub>), 108.08(=C= of Dde), 79.47 [COO-C(Me)<sub>3</sub>], 65.83 (O-CH<sub>2</sub>-CH=), 60.99 (-CH<sub>2</sub>- of Dde), 53.07, 47.83, 43.40, 40.55, 32.94, 30.35, [28.96, 28.68, 28.52, -COO-C(CH<sub>3</sub>)<sub>3</sub>], 23.35, 87, 18.18 (CH<sub>3</sub> of Dde).

C.H.N analyse for C<sub>28</sub>H<sub>43</sub>N<sub>3</sub>O<sub>6</sub>·H<sub>2</sub>O: calc. C 61.03, H 8.86, N 8.21; found C 61.12, H 8.78, N 8.13.

**EXAMPLE 18****N-(2-Boc-aminoethyl)-N-(Thymine-1-ylacetyl)-Lysine-N'-(Dde) O-Allyl (18)***Preparation: Procedure (c), light yellow crystalline. Yield = 87%*

Mp = 84-86; MS (FAB)  $m/z$  660 (M+H);  $^1\text{H-NMR}$ , ( $\text{CDCl}_3$ ):  $\delta$  13.4 (s, 1H, -HN-Dde), 9.56 (s, 1H, aromatic proton of Thymine), 6.99 (s, 1H, -NH- of Thymine), 5.97-5.84 (m, 1H, -CH=CH<sub>2</sub>), 5.66 (s, 1H, boc-NH-), 5.35 (dt, 1H,  $J_{\text{trans}}=17.28$ , 1.46 Hz), 5.25 (dd, 1H,  $J_{\text{cis}}=10.55$ , 1.17 Hz), 4.7-4.4 (overlapping m and s, 4H, O-CH<sub>2</sub>-CH=, CO-CH<sub>2</sub>-Thymine), 4.2 (t, 1H), 3.7 (m, 1H), 3.44-3.27 (overlapping m, 5H), 2.55 (s, 3H, -HN-C-CH<sub>3</sub>), 2.37 (s, 4H, 2 x-CH<sub>2</sub>- of Dde), 2.05 (m, 4H), 1.89 (s, 3H, CH<sub>3</sub> of Thymine), 1.7 (m, 2H), 1.45 (s, 9H, boc), 1.03 (s, 6H, 2x-CH<sub>3</sub> of Dde);  $^{13}\text{C-NMR}$  ( $\text{CDCl}_3$ ):  $\delta$  = 197.81 (-CO- of Dde), 173.54, 170.33, 167.50, 163.94, 155.89 (-NH-COO), 150.74, 140.91, 131.26 (-CH=CH<sub>2</sub>), 119.26 (-CH=CH<sub>2</sub>), 110.41, 107.80 (=C= of Dde), 79.85 [-COO-C(Me)<sub>3</sub>], 66.25 (-CH<sub>2</sub>-CH=), 60.27 (-CH<sub>2</sub>- of Dde), 52.69, 48.30, 47.66, 42.93, 39.11, 30.01, [28.31, 28.13, 28.01, -COO-C(CH<sub>3</sub>)<sub>3</sub>], 23.36, 17.89 (CH<sub>3</sub> of Dde), 12.29 (CH<sub>3</sub> of Thymine).

C.H.N analyse for C<sub>33</sub>H<sub>49</sub>N<sub>5</sub>O<sub>9</sub>: calc. C 60.08, H 7.49, N 10.61; found C 59.70, H 7.49, N 10.50.

**EXAMPLE 19****N-(2-Boc-aminoethyl)-N-(Thymine-1-ylacetyl)-Lysine-O-Allyl (19)**

*Preparation:* 1.4 g (2.1 mmol) **18** was dissolved in a solution of 18 ml allyl alcohol and 2 ml hydrazin hydrate and stirred for 15 min. 200 ml DMF was added and the reaction mixture

was evaporated under high vacuum. 50 ml Water was added to the remaining oil and extracted with diethylether (3 x 50 ml). The water phase was freeze dried and 0.8 g (1.6 mmol) titled compound was obtained as slightly yellow crystalline, which was used for the next step without further purification (yield = 76%). A sample of product was purified by HPLC and collected as TFA salt.

Mp = 81-83; MS (FAB)  $m/z$  496 (M+H);  $^1\text{H-NMR}$ , ( $\text{CD}_3\text{OD}$ ):  $\delta$  7.34 (s, aromatic proton of Thymine), 6.03-5.94 (12 line m, 1H,  $-\text{CH}=\text{CH}_2$ ), 5.37(ddd, 1H,  $J_{\text{trans}}=17.21, 3.11, 1.46$  Hz), 5.27 (dd, 1H,  $J_{\text{cis}}=10.44, 2.56, 1.28$  Hz), 4.8-4.6 (ddt, 2H,  $J=32, 5.67, 1.28$  Hz,  $\text{O}-\text{CH}_2-\text{CH}=\text{}$ ), 4.78 (s, 2H,  $\text{CO}-\text{CH}_2-\text{Thymine}$ ), 4.37 (dd, 1H,  $J=8.8, 5.86$  Hz,  $^a\text{CH}$ ), 3.7 (m, 1H,  $\text{boc-HN}-\text{CH}_2-\text{CHH}'$ ), 3.4 [overlapping m and d ( $J=0.74$ ), 3H,  $\text{boc-HN}-\text{CH}_2-\text{CHH}'$ ,  $-\text{NH}_2$ ], 3.23 (m, 1H), 3.03 (m, 1H), 3.97 (t, 2H,  $j=7.51, -\text{CH}_2-\text{NH}_2$ ), 2.2-2 (dm, 2H,  $^b\text{CH}_2$ ), 1.9 (s, 3H,  $\text{CH}_3$  of Thymine) 1.7 (m, 2H), 1.49 [overlapping s and m, 11H,  $\text{boc}(9\text{H}), -\text{CH}_2-\text{CH}_2-\text{NH}_2$ ];  $^{13}\text{C-NMR}$  ( $\text{CD}_3\text{OD}$ ): (major rotamer)  $\delta$  171.79, 169.81, 166.98, 158.50 ( $-\text{NH}-\text{COO}-t\text{-butyl}$ ), 153.12, 143.74, 133.44 ( $-\text{CH}=\text{CH}_2$ ), 119.03 ( $-\text{CH}=\text{CH}_2$ ), 116.38, 111.11, 80.74 [ $-\text{COO}-\text{C}(\text{Me})_3$ ], 67.21 ( $-\text{CH}_2-\text{CH}=\text{}$ ), 61.94, 50.21, 40.61, 29.61, 28.81, 28.37, 27.83, 24.30, 12.32 ( $\text{CH}_3$  of Thymine). C.H.N analyse for  $\text{C}_{23}\text{H}_{37}\text{N}_5\text{O}_7 \cdot 2 \text{CF}_3\text{COOH}$ : calc. C 44.82, H 5.43, N 9.68; found C 44.61, H 5.66, N 9.92.

#### EXAMPLE 20

#### ***N*-boc-*N*-(Thymine-1-ylacetyl)Lysine-[*N*-*N*-bis(2,3,4,5-tetra-*O*-acetyl Galactose-1-yl)-ethyl]-*O*-Allyl (20)**

*Preparation:* 300mg (0.6 mmol) **19** was dissolved in 10 ml methanol, 500 mg (1.33 mmol, 2.2 eq) (**M5**) was added and stirred for 10 min. 80  $\mu\text{ml}$  (1.3 mmol) Acetic acid and 0.6g (10 mmol) natrium cyanoborohydrate was added sequentially. After 15 min the same amounts of acetic acid and  $\text{NaCNBH}_3$  were added and the reaction mixture was stirred for further 30 min at room temperature. Volatile was removed under vacuum and remaining dissolved in 100 ml ethylacetate and extracted with  $\text{NaHCO}_3$  (2 x 50ml) saturated solution and 50 ml brine respectively. Organic phase was dried over magnesium sulphate and evaporated to dryness under vacuum. The residue was purified on silica gel column eluting ethylacetate-methanol 10:0 to 10:0.5. 440 mg (0.36 mmol) titled compound was obtained as white crystalline. Yield = 60%.

Mp = 106-108; MS (FAB)  $m/z$  1212 (M+H);  $^1\text{H-NMR}$ , ( $\text{CDCl}_3$ ): resolved signals:  $\delta$  7 (s, 1H, aromatic proton of Thymine), 5.9 (m, 1H,  $-\text{CH}=\text{CH}_2$ ), 5.5 (br.s, 1H,  $-\text{NH}-\text{boc}$ ), 5.37 (d, 2H,  $J=2.74$  Hz,  $2x^4\text{CH-sugar}$ ), 5.3 (dd, 1H,  $J_{\text{trans}}=17.2, 1.28$  Hz), 5.2 (overlapping, 5H,  $-\text{CH}=\text{CHH}'$ ,  $2x^3\text{CH-}$  and  $2x^2\text{CH-}$  of sugar), 4.64 (s, 2H), 4.58 (d, 2H,  $J=5.86$  Hz,  $\text{O}-\text{CH}_2-\text{CH}=\text{}$ ), 4.29 (br.d,

2H,  $J=10.07$ ), 4.15 (br., 2H), 4 (br.d 2H,  $J=8.4$ ), 2.2-2 (4s, 24H, CO-CH<sub>3</sub> of sugar), 1.8 (s, 3H, CH<sub>3</sub> of Thymine), 1.4 (overlapping s, 11H, 9H of boc, 2H of lysine side chain); <sup>13</sup>C-NMR (CDCl<sub>3</sub>):  $\delta$  170.75, 169.97, 169.65, 169.59, 167.58, 164.13, 156.00, 151.06, 141.38, 131.24, 119.06, 110.31, 79.65, 68.20, 67.44, 66.65, 66.18, 60.75, 60.18, 50.01, 39.12, 28.25, 20.66, 20.61, 20.48, 12.00.

HRMS (M+H)<sup>+</sup>, calcd (found) for C<sub>55</sub>H<sub>82</sub>N<sub>5</sub>O<sub>25</sub> are 1212.5299 (1212.5360).

#### EXAMPLE 21

##### ***N*-boc-*N*-(Thymine-1-ylacetyl)Lysine-[*N*-*N*-bis(2,3,4-tri-*O*-acetyl Fucose-1-yl)-ethyl]-*O*-Allyl (21)**

*Preparation:* The Procedure is the same as described above for the synthesis of 20. yield = 63%.

Mp = ; MS (FAB)  $m/z$  1096 (M+H); <sup>1</sup>H-NMR, (CDCl<sub>3</sub>): resolved signals:  $\delta$  7 (s, 1H, aromatic proton of Thymine), 5.9 (m, 1H, -CH=CH<sub>2</sub>), 5.6 (br.s, 1H, -NH-boc), 5.3 (s, 1H), 5.2 (overlapping, 4H, -CH=CHH', sugar), 4.58 (s, 2H, O-CH<sub>2</sub>-CH=), 4.2 (br.s, 1H), 4 (dd, 2H,  $J=7.3$ , 0.8 Hz), 3.4-3.2 (br. 2H), 2.9 (br.s, 1H), 2.8 (br.s, 1H), 2 (3s, 18H, CO-CH<sub>3</sub> of sugar), 1.8 (s, 3H, CH<sub>3</sub> of Thymine), 1.4 (s, 9H, boc), 1.2 (dd, 2H,  $J=7.3$ , 1.2 of lysine side chain), 1.1 (d, 6H,  $J=3.8$ , methyl group of fucose); <sup>13</sup>C-NMR (CDCl<sub>3</sub>):  $\delta$  170.87, 170.45, 170.17, 169.95, 167.21, 164.46, 155.90, 151.11, 141.38, 131.18, 130.89, 119.34, 118.79, 110.26, 80.08, 72.51, 70.39, 69.75, 69.50, 68.07, 67.86, 66.05, 60.80, 60.08, 53.51, 50.20, 48.69, 48.01, 38.89, 36.44, 31.28, 28.09, 23.63, 21.65, 20.73, 20.60, 20.48, 20.38, 16.00, 15.47, 13.87, 11.94.

HRMS (M+H)<sup>+</sup>, calcd (found) for C<sub>51</sub>H<sub>78</sub>N<sub>5</sub>O<sub>21</sub> are 1096.5189(1096.5192).

#### EXAMPLE 22

##### ***N*-boc-*N*-(Thymine-1-ylacetyl)Lysine-[*N*-(2,3,4,5-tetra-*O*-acetyl Galactose-1-yl)-acetyl]-*O*-Allyl (22)**

1.03 g (2.64 mmol, 1.1 eq) (M6), 470 mg (2.88 mmol, 1.2 eq) DhbtOH, and 740mg (3.6 mmol, 1.5 eq) DCC were mixed in 20 ml DMF and stirred for 30 min under Nitrogen. A solution of 1.2 g (2.4 mmol) 19 in 10 ml DMF was added and the reaction mixture was stirred for further 3h. Volatiles were removed under vacuum and the residue dissolved in 200ml ethylacetate. Insoluble DCU was filtered off and the filtrate was extracted with NaHCO<sub>3</sub> saturated solution (2 x 100 ml) and brine (100 ml) respectively. Organic phase was dried over magnesium sulphate and evaporated to dryness under vacuum. The remaining was purified on silica gel column eluting ethylacetate. 1.1 g (1.2 mmol) title compound was obtained as white crystalline. Yield = 61%.

Mp = 82-84; MS (FAB)  $m/z$  868 (M+H);  $^1\text{H-NMR}$ , ( $\text{CDCl}_3$ ): resolved signals:  $\delta$  9.39 (aromatic proton of Thymine), 6.94 (s, 1H, -NH- of Thymine), 6.64 (s, 1H, -NH-CO-sugar), 5.92-5.85 (m, H,  $-\underline{\text{C}}\text{H}=\text{CH}_2$ ), 5.42 (overlapping d, 2H,  $J=2.75$ ,  $^4\text{CH}$ -sugar, -NH-boc), 5.32 (dd, 1H,  $J_{\text{trans}}=17.2$ , 1.46 Hz), 5.25 (overlapping, 3H,  $-\text{CH}=\text{CH}\underline{\text{H}}$ ,  $^3\text{CH}$ - and  $^2\text{CH}$ - of sugar), 4.7 (m, 1H), 4.6 (d, 2H,  $J=5.86$ ), 4.5 (m, 1H), 4.3-4.1 (m, 5H), 3.6 (m, 1H), 3.35 (m, 3H), 3.2 (br., 2H), 2.6 (m, 1H), 2.45 (m, 1H), 2.15-2 (3s, 12H,  $\text{CO-CH}_3$  of sugar), 1.9 (s, 3H,  $\text{CH}_3$  of Thymine), 1.6 (br., 2H), 1.4 (overlapping s, 11H, 9H of boc, 2H of lysine side chain);  $^{13}\text{C-NMR}$  ( $\text{CDCl}_3$ ):  $\delta$  170.56, 169.85, 169.77, 169.44, 167.47, 163.93, 155.82, 151.40, 140.89, 131.23 ( $-\underline{\text{C}}\text{H}=\text{CH}_2$ ), 119.09 ( $-\text{CH}=\underline{\text{C}}\text{H}_2$ ), 110.93, 79.86 [ $-\text{O}-\underline{\text{C}}(\text{Me})_3$ ], 69.23, 68.87, 68.23, 67.60, 66.82, 66.12, 60.85, 59.64, 48.88, 47.13, 38.99, 38.59, 34.74, 28.28, 28.04, 27.58, 22.79, 20.61, 20.51, 12.20 ( $\text{CH}_3$  of Thymine)

C.H.N analyse for  $\text{C}_{39}\text{H}_{57}\text{N}_5\text{O}_{17} \cdot 1/2 \text{H}_2\text{O}$ : calc. C 53.42, H 6.67, N 7.99; found C 53.25, H 6.71, N 7.44.

#### 15 EXAMPLE 23

***N*-boc-*N*-(Thymine-1-ylacetyl)Lysine-[*N*-(2,3,4,5-tetra-*O*-acetyl-Mannose-1-yl)-acetyl]-*O*-Allyl (23)**

*Preparation:* The Procedure is the same as described above for the synthesis of **22**. Yield = 58%.

20 Mp = ; MS (FAB)  $m/z$  868 (M+H);  $^1\text{H-NMR}$ , ( $\text{CDCl}_3$ ): resolved signals:  $\delta$  9.35 (aromatic proton of Thymine), 6.94 (s, 1H, -NH- of Thymine), 6.78 (s, 1H, -NH-CO-sugar), 5.8 (m, 1H,  $-\underline{\text{C}}\text{H}=\text{CH}_2$ ), 5.45 (s, 1H, -NH-boc), 5.35 (dd, 1H,  $J_{\text{trans}}=17$ , 1.2 Hz), 5.25 (m, 2H of sugar), 4.6 (d, 2H,  $J=5.5$ ), 4.5 (s, 1H), 4.3 (dd, 1H,  $J=12.1$ , 5.7 Hz), 4.2 (dd, 1H,  $J=12.1$ , 8.2), 4.1 (m, 1H), 3.6 (m, 1H), 3.35 (m, 5H), 2.6 (m, 2H), 2.15-2 (4s, 12H,  $\text{CO-CH}_3$  of sugar), 1.9 (s, 3H,  $\text{CH}_3$  of Thymine), 1.6 (br., 2H), 1.4 (overlapping s, 11H, 9H of boc, 2H of lysine side chain);  $^{13}\text{C-NMR}$  ( $\text{CDCl}_3$ ):  $\delta$  170.73, 170.19, 169.97, 169.48, 169.28, 167.36, 163.82, 156.56, 151.39, 143.55, 141.16, 131.21 ( $-\underline{\text{C}}\text{H}=\text{CH}_2$ ), 127.70, 127.01, 124.81, 119.93, 119.21 ( $-\text{CH}=\underline{\text{C}}\text{H}_2$ ), 110.87, 71.56, 70.33, 69.35, 68.11, 67.26, 66.67, 66.24, 61.73, 59.90, 47.06, 39.51, 38.79, 36.95, 28.14, 27.62, 22.85, 20.66, 12.18 ( $\text{CH}_3$  of Thymine).

30

#### EXAMPLE 24

***N*-boc-*N*-(Thymine-1-ylacetyl)Lysine-[*N*-(2,3,4,5-tetra-*O*-acetyl fucose-1-yl)-acetyl]-*O*-Allyl (24)**

*Preparation:* The Procedure is the same as described above for the synthesis of **22**. Yield = 65%.

35

Mp= ; MS (FAB)  $m/z$  810 (M+H);  $^1\text{H-NMR}$ , ( $\text{CDCl}_3$ ): resolved signals:  $\delta$  9.64 (aromatic proton of Thymine), 6.94 (s, 1H, -NH- of Thymine), 6.88 (s, 1H, -NH-CO-sugar), 6-5.8 (m, H, -CH=CH<sub>2</sub>), 5.5 (s, 1H, -NH-boc), 5.37 (d, 1H,  $J=3.8$ ) 5.3 (overlapping, 2H, -CH=CHH'), 5.16 (m, 1H of sugar), 4.7 (d, 1H,  $J=15.5$ ), 4.65 (m, 1H), 4.6 (d, 2H,  $J=5.8$ ), 4.3 (m, 1H), 4.1 (m, 1H), 3.9 (m, 1H), 3.8 (m, 1H), 3.5-3.1 (m, 5H), 2.9 (dd, 1H,  $J=23.1, 2.3\text{Hz}$ ), 2.7 (m, 1H), 2.4-2.2 (m, 2H), 2.15-2 (3s, 9H, CO-CH<sub>3</sub> of sugar), 1.9 (s, 3H, CH<sub>3</sub> of Thymine), 1.4 (s, 9H of boc), 1.2 (m, 2H, of lysine side chain);  $^{13}\text{C-NMR}$  ( $\text{CDCl}_3$ ):  $\delta$  170.42, 169.95, 169.74, 169.68, 169.55, 167.13, 163.86, 162.37, 155.81, 151.53, 140.81, 131.25 (-CH=CH<sub>2</sub>), 119.08 (-CH=CH<sub>2</sub>), 111.01, 79.83 [-O-C(Me)<sub>3</sub>], 69.36, 68.43, 68.23, 67.95, 66.16, 61.64, 60.24, 48.95, 38.83, 37.41, 36.34, 35.68, 33.78, 28.72, 28.34, 27.33, 27.46, 24.80, 22.67, 20.91, 20.69, 20.62, 14.63, 14.05, 12.30 (CH<sub>3</sub> of Thymine).

#### EXAMPLE 25

***N*-Fmoc-*N*-(Thymine-1-ylacetyl)Lisine-[*N*-(2,3,4,5-tetra-*O*-acetyl Galactose-1-yl)-acetyl]-*O*-Allyl (25)**

*Preparation: procedure (d). Yield = 91%.*

Mp = 93-95; MS (FAB)  $m/z$  990 (M+H);  $^1\text{H-NMR}$ , ( $\text{CDCl}_3$ ):  $\delta$  9.33 (aromatic proton of Thymine), 7.6-7.1 (8H, florenyl aromatics), 6.64 (s, 1H, -NH- of Thymine), 6.51 (s, 1H, -NH-CO-sugar), 5.8-5.6 (overlapping m, 2H, -CH=CH<sub>2</sub>, -NH-fmoc), 5.27 (s, 1H, <sup>4</sup>CH-sugar), 5.15 (d, 1H,  $J_{\text{trans}}=16.66\text{ Hz}$ ), 5.07 (overlapping, 3H, -CH=CHH', <sup>3</sup>CH and <sup>2</sup>CH of sugar), 4.55 (br.d, 1H,  $J=3.84$ ), 4.45 (d, 2H,  $J=5.86\text{ Hz}$ , O-CH<sub>2</sub>-CH=), 4.28 (overlapping s, 3H), 4.2 (d, 1H,  $J=7.14$ ), 4.1-4 (m, 5H), 3.42 (m, 1H), 3.29-3.26 (overlapping s, 3H), 3.04 (br.s, 2H), 2.46 (dd, 1H,  $J=14.83, 9.15\text{ Hz}$ ), 2.26 (m, 1H), 2-1.8 (4s, 12H, -CO-CH<sub>3</sub> of sugar), 1.68 (s, 3H, CH<sub>3</sub> of Thymine), 1.4 (m, 1H), 1.2 (m, 3H);  $^{13}\text{C-NMR}$  ( $\text{CDCl}_3$ ):  $\delta$  170.56, 170.46, 169.80, 169.72, 169.41, 169.36, 167.51, 163.98, 156.48, 151.29, 143.47, 143.44, 141.07, 131.18, 127.60, 127.49, 126.89, 124.71, 119.83, 119.03, 110.59, 77.41, 76.90, 76.39, 69.21, 68.75, 68.17, 67.64, 67.52, 66.73, 66.48, 66.06, 60.77, 59.54, 48.68, 46.97, 39.52, 38.62, 34.66, 28.16, 27.69, 22.87, 20.52, 20.43, 12.08.

HRMS (M+H)<sup>+</sup>, calcd (found) for C<sub>49</sub>H<sub>80</sub>N<sub>5</sub>O<sub>17</sub> are 990.3984(990.3940).

#### EXAMPLE 26

***N*-Fmoc-*N*-(Thymine-1-ylacetyl)Lysine-[*N*-(2,3,4,5-tetra-*O*-acetyl mannose-1-yl)-acetyl]-*O*-Allyl (26)**

*Preparation: procedure (d). Yield = 88%.*



Mp= ; MS (FAB)  $m/z$  990 (M+H);  $^1\text{H-NMR}$ , ( $\text{CDCl}_3$ ):  $\delta$  9.24 (aromatic proton of Thymine), 7.7-7.2 (8H, florenyl aromatics), 6.71 (overlapping-s, 2H, -NH- of Thymine, -NH-CO-sugar), 5.8 (overlapping-m, 2H, -CH=CH<sub>2</sub>, -NH-fmoc), 5.28 (s, 1H, sugar), 5.2(m, 2H, -CH=CHH'), 5.06 (t, 2H, J=6.7), 4.55 (d, 2H, J=4.1), 4.4 (3, 2H, O-CH<sub>2</sub>-CH=), 4.3 (m, 1H), 4.2 (m, 1H), 4.1 (m, 1H), 3.6 (m, 1H), 3.4-3.1 (m, 3H), 2.5 (m, 2H), 2 (s, 12H, -CO-CH<sub>3</sub> of sugar), 1.78 (s, 3H, CH<sub>3</sub> of Thymine), 1.5 (m, 1H), 1.2 (m, 3H);  $^{13}\text{C-NMR}$  ( $\text{CDCl}_3$ ):  $\delta$  170.72, 169.54, 165.72, 163.81, 151.40, 143.57, 141.17, 136.86, 131.22, 127.70, 127.01, 124.89, 119.93, 119.22, 110.10, 81.47, 67.27, 66.65, 66.24, 61.74, 47.07, 20.65, 20.43, 12.18.  
HRMS (M+H)<sup>+</sup>, calcd (found) for C<sub>48</sub>H<sub>60</sub>N<sub>5</sub>O<sub>17</sub> are 990.3984(990.3987).

#### EXAMPLE 27

##### ***N*-Fmoc-*N*-(Thymine-1-ylacetyl)Lysine-[*N*-(2,3,4-tri-*O*-acetyl fucose-1-yl)-acetyl]-*O*-Allyl (27)**

*Preparation: procedure (d).* Yield = 89% **3**

Mp= ; MS (FAB)  $m/z$  932 (M+H);  $^1\text{H-NMR}$ , ( $\text{CDCl}_3$ ):  $\delta$  9.85 (aromatic proton of Thymine), 7.7-7.2 (8H, florenyl aromatics), 7 (s, 1H, -NH- of Thymine), 6.8 (s, 1H, -NH-CO-sugar), 6 (s, 1H, -NH-fmoc), 5.9 (m, 1H, -CH=CH<sub>2</sub>), 5.28 (s, 1H, sugar), 5.2 (m, 2H, -CH=CHH'), 5.1 (s, 1H), 4.6 (overlapping-s, 3H), 4.4 (m, 2H), 4.25 (br.s, 1H), 4.2(m, 1H), 2 (s, 9H, -CO-CH<sub>3</sub> of sugar), 1.8 (s, 3H, CH<sub>3</sub> of Thymine), 1.5-1.2 (m, 4H of Lysine side chain);  $^{13}\text{C-NMR}$  ( $\text{CDCl}_3$ ):  $\delta$  170.99, 170.46, 170.16, 169.98, 169.79, 169.62, 167.11, 163.90, 156.50, 151.60, 143.65, 143.31, 141.11, 131.22, 130.90, 127.65, 127.02, 126.96, 124.87, 119.88, 119.13, 110.87, 69.05, 68.19, 67.70, 66.60, 66.20, 61.55, 60.23, 48.77, 46.99, 39.32, 37.69, 35.20, 28.77, 27.46, 22.74, 20.89, 20.61, 20.56, 20.47, 14.72, 14.02, 12.15.  
HRMS (M+H)<sup>+</sup>, calcd (found) for C<sub>47</sub>H<sub>58</sub>N<sub>5</sub>O<sub>15</sub> are 932.3929(932.3945).

#### EXAMPLE 28

##### ***N*-(2-Fmoc-aminoethyl)- Asp(tBu)-Allyl (30)**

To 4.5 g (10 mmol) **28** [prepared by procedure (a)] was added 50 ml of a solution of 20% piperidine in DCM. After 30 min, 100 ml toluene was added and volatile were removed under vacuum. The residue was purified with a short silica gel column eluting ethylacetate-methanole 10:0 to 10:1. 1.8 g (7.9 mmol) **29** was obtained as slightly yellow oil. This oil was used for *procedure (b)*. 1.7 g (3.4 mmol) title compound was obtained as colourless oil. Overall yield = 34%.

[ $\alpha$ ]<sub>D</sub><sup>22</sup> = (c=1, methanol); MS (FAB) *m/z* 495 (M+H); <sup>1</sup>H-NMR, (CDCl<sub>3</sub>):  $\delta$  7.7-7.2 (8H, Florenyl aromatic protons), 5.96-5.84 (16 line m, 1H, -CH=CH<sub>2</sub>), 5.56 (br.s, 1H, fmoc-HN-), 5.35 (dt, 1H, *J*<sub>trans</sub>=15.9, 1.46 Hz), 5.25 (dt, 1H, *J*<sub>cis</sub>=10.5, 1.28 Hz), 4.6 (dd, 2H, *J*=5.9, 1.1 Hz, O-CH<sub>2</sub>-CH=), 4.4 (d, 2H, *J*=7 Hz), 4.2 (t, 1H, *J*=7 Hz), 3.6 (t, 1H, *J*=7.3 Hz, <sup>o</sup>CH), 3.3 (m, 2H, <sup>p</sup>CH), 2.9 (m, 1H), 2.7 (m, 2H), 2.6 (m, 1H), 2.3 (br.s, 1H), 1.4 (s, 9H, *t*-butyl) <sup>13</sup>C-NMR (CDCl<sub>3</sub>):  $\delta$  173.02 (COO-Allyl), 169.89, 156.37, 143.84, 141.06, 131.47(-CH=CH<sub>2</sub>), 127.41, 126.81, 124.95, 119.71, 118.68(-CH=CH<sub>2</sub>), 81.17[-O-C(Me)<sub>3</sub>], 66.5 (-CH<sub>2</sub>-CH=), 65.62 ( $\beta$  carbon), 57.12 ( $\alpha$  carbon), 47.05, 40.34, 38.97, 27.85 [-COO-C(CH<sub>3</sub>)<sub>3</sub>]. HRMS (M+H)<sup>+</sup>, calcd (found) for C<sub>28</sub>H<sub>35</sub>N<sub>2</sub>O<sub>8</sub> are 495.2495(495.2480).

10

## EXAMPLE 29

***N*-(2-fmoc-aminoethyl)-*N*-(Thymine-1-ylacetyl)-Asp-(*t*Bu)-O-Allyl (31)**

*Preparation: procedure (c).* Yield = 81%.

Mp = ; [ $\alpha$ ]<sub>D</sub><sup>22</sup> = (c=0.25, methanol); MS (FAB) *m/z* 661 (M+H); <sup>1</sup>H-NMR, (CDCl<sub>3</sub>):  $\delta$  9.15 (s, 1H, Thymine aromatic), 7.75-7.26 (8H, Florenyl aromatic protons), 6.75 (s, 1H, Thymine-NH-), 5.9 (s, 1H, fmoc-HN-), 5.8 (m, 1H, -CH=CH<sub>2</sub>), 5.3 (d, 1H, *J*<sub>trans</sub>=17.3 Hz), 5.2 (d, 1H, *J*<sub>cis</sub>=10.5 Hz), 4.6 (m, 3H, O-CH<sub>2</sub>-CH=, <sup>o</sup>CH), 4.45 (s, 2H, -CO-CH<sub>2</sub>-Thymine), 4.2-4.1 (m, 3H), 3.4-3.4 (m, 2H), 3.4-3.2 (m, 2H), 2.9-2.7 (m, 2H), 1.8 (s, 3H, Thymine-CH<sub>3</sub>), 1.4 (s, 9H, -COOtBu); <sup>13</sup>C-NMR (CDCl<sub>3</sub>):  $\delta$  170.49, 169.36, 167.05, 164.06, 156.52, 150.62, 143.63, 143.52, 141.09, 140.85, 131.10(-CH=CH<sub>2</sub>), [(130.71, 127.57, 126.92, 124.83, 119.79(florenyl)], 119.24(-CH=CH<sub>2</sub>), 110.32, 82.27, 81.41, 66.84, 66.51, 60.24(-CH<sub>2</sub>-CH=), 58.08, 48.85, 47.82, 47.10, 39.15, 35.41, 27.85, 14.04, 12.19(Thymine-CH<sub>3</sub>). HRMS (M+H)<sup>+</sup>, calcd (found) for C<sub>35</sub>H<sub>41</sub>N<sub>4</sub>O<sub>8</sub> are 661.2874(661.2881).

## 25 EXAMPLE 30

***N*-(2-fmoc-aminoethyl)-*N*-(Thymine-1-ylacetyl)-Asp-O-Allyl (32)**

*Preparation:* To 3.3 g (5 mmol) **31** was added a solution of 95% TFA 5% TES. The reaction mixture was stirred until total conversion of starting material according to TLC. TFA solution was co-evaporated with DCM under vacuum. Titled compound was obtained as slightly yellow crystalline which was used for the next step without further purification (Yield = 62%); Mp =

30

[ $\alpha$ ]<sub>D</sub><sup>22</sup> = (c=0.25, methanol); MS (FAB)=605 *m/z* (M+H); <sup>1</sup>H-NMR, (DMSO):  $\delta$  11.3 (s, 1H, Thymine aromatic), 7.91-7.22 (8H, Florenyl aromatic protons, 1H, Thymine-NH-), 5.8 (m, 1H, -CH=CH<sub>2</sub>), 5.3 (dd, 1H, *J*<sub>trans</sub>=17.3, 1.76 Hz), 5.1 (dd, 1H, *J*<sub>cis</sub>=10.55, 1.47 Hz), 4.6 (d, 2H, *J*=7 Hz, O-CH<sub>2</sub>-CH=), 4.5 (s, 2H, -CO-CH<sub>2</sub>-Thymine), 4.4 (t, 1H, *J*=6.4, <sup>o</sup>CH), 4.3 (d, 2H, *J*=6.8, florenyl-CH<sub>2</sub>-O-), 4.2 (t, 1H, *J*=6.8, florenyl-CH<sub>2</sub>-O-), 3.5 (m, 2H), 3.4 (m, 2H), 3.2

35

(dd, 2H,  $J=17, 7.3$  Hz), 1.7 (s, 3H, Thymine-CH<sub>3</sub>); <sup>13</sup>C-NMR (CD<sub>3</sub>OD):  $\delta$  174.50, 170.90, 169.54, 166.94, 158.90, 152.80, 145.30, 143.64, 142.57, 142.16, 133.25, 131.09 (-CH=CH<sub>2</sub>), [(130.44, 128.83, 128.21, 127.13, 126.20, 121.00(florenyl))], 118.96 (-CH=CH<sub>2</sub>), 110.89, 67.80, 67.38, 61.59, 59.96(-CH<sub>2</sub>-CH=), 40.47, 35.32, 14.53, 12.34 (Thymine-CH<sub>3</sub>).

5 HRMS (M+H)<sup>+</sup>, calcd (found) for C<sub>31</sub>H<sub>33</sub>N<sub>4</sub>O<sub>9</sub> are 605.2224(605.2248).

#### EXAMPLE 31

##### **fmoc-N-(Thymine-1-ylacetyl)Asp-[O-(2,3,4,5-tetra-O-acetyl Galactose-1-yl)-amide]-O-Allyl (33)**

10 *Preparation: procedure (e).* Yield = 64%.

Mp = ;  $[\alpha]_D^{22}$  (c=0.5, methanol); MS (FAB)  $m/z$  934 (M+H); <sup>1</sup>H-NMR, (CDCl<sub>3</sub>):  $\delta$  9.5 (s, 1H, Thymine aromatic), 7.7-7.2 (m, 8H, Florenyl aromatic protons), 6.6 (s, 1H, Thymine-NH-), 5.9 (overlapping m, 2H, -CH=CH<sub>2</sub>, fmoc-HN-), 5.3 (d, 1H,  $J=1.6$ Hz), 5.25-5.15 (m, 3H), 5.05 (d, 2H,  $J=5.3$  Hz), 4.6 (m, 2H), 4.4 (m, 2H), 4.1 (m, 1H), 4 (m, 3H), 3.6-3.2 (m, 5H), 2.8 (m, 1H), 2.1-1.9 (4s, 12H, sugar acetyl protons), 1.8 (s, 3H, Thymine-CH<sub>3</sub>); <sup>13</sup>C-NMR (CDCl<sub>3</sub>):  $\delta$  171.21, 170.22, 169.84, 169.76, 169.58, 166.56, 164.17, 156.81, 156.57, 151.39, 143.68, 143.45, 141.17, 140.94, 131.20(-CH=CH<sub>2</sub>), 127.71, 127.04, 124.89, 119.92, 119.16 (-CH=CH<sub>2</sub>), 110.60, 80.39, 78.26, 72.13, 71.36, 67.86, 67.19, 66.81, 66.55, 61.44, 61.06, 57.81, 49.22, 48.92, 48.77, 47.16, 39.19, 36.00, 33.69, 25.47, 24.79, 20.71, 20.61, 20.57, 20.46 (sugar-CO-CH<sub>3</sub> groups), 12.26 (Thymine-CH<sub>3</sub>).

C.H.N analyse for C<sub>45</sub>H<sub>51</sub>N<sub>5</sub>O<sub>17</sub>, H<sub>2</sub>O: calc. C 56.78, H 5.61, N 7.36; found C 56.87, H 5.94, N 7.57.

#### EXAMPLE 32

##### **Fmoc-N-(Thymine-1-ylacetyl)Asp-[O-(2,3,4,5-tetra-O-acetyl mannose-1-yl)-amide]-O-Allyl (34)**

*Preparation: procedure (e).* Yield = 60%.

Mp = ;  $[\alpha]_D^{22}$  (c=0.5, methanol); MS (FAB)  $m/z$  934(M+H); <sup>1</sup>H-NMR, (CDCl<sub>3</sub>):  $\delta$  9.5 (s, 1H, Thymine aromatic), 7.7-7.2 (m, 8H, Florenyl aromatic protons), 6.7 (s, 1H, Thymine-NH-), 6 (br.s, 1H, fmoc-HN-), 5.9 (m, 1H, -CH=CH<sub>2</sub>), 5.5 (d, 1H,  $J=8.8$ Hz), 5.4 (d, 1H,  $J=8.8$  Hz), 5.3-5 (m, 5H), 4.6 (m, 2H), 4.4 (m, 2H), 4.2 (m, 3H), 4.1 (m, 2H), 3.8-3.4 (m, 5H), 3.1 (m, 1H), 2.9 (m, 1H), 2.2-1.9 (4s, 12H, sugar acetyl protons), 1.8 (s, 3H, Thymine-CH<sub>3</sub>); <sup>13</sup>C-NMR (CDCl<sub>3</sub>). Mp = ;  $[\alpha]_D^{22}$  (c=0.5, methanol); MS (FAB)  $m/z$  934 (M+H); <sup>1</sup>H-NMR, (CDCl<sub>3</sub>) <sup>13</sup>C-NMR (CDCl<sub>3</sub>):  $\delta$  171.13, 170.95, 170.49, 170.30, 169.93, 169.85, 164.62, 157.14, 156.95, 151.54, 144.08, 141.49, 131.57(-CH=CH<sub>2</sub>), 128.04, 127.35, 125.24, 120.27, 119.56 (-

CH=CH<sub>2</sub>), 110.77, 74.43, 71.83, 69.87, 67.00, 66.87, 66.05, 65.62, 62.61, 58.53, 49.56, 49.34, 47.44, 39.47, 34.13, 25.83, 25.16, 21.21, 21.01, 20.95, 20.84 (sugar-CO-CH<sub>3</sub> groups), 12.05 (Thymine-CH<sub>3</sub>).

HRMS (M+H)<sup>+</sup>, calcd (found) for C<sub>46</sub>H<sub>52</sub>N<sub>5</sub>O<sub>17</sub> is 934.3358 (934.3342).

5

#### EXAMPLE 33

##### **Fmoc-N-(Thymine-1-ylacetyl)Asp-[O-(2,3,4-tri-O-acetyl fucose-1-yl)-amide]-O-Allyl (35)**

*Preparation: procedure (e).* Yield = 77%.

Mp = ;  $[\alpha]_D^{22}$  (c=0.5, methanol); MS (FAB) *m/z* 876 (M+H); <sup>1</sup>H-NMR, (CDCl<sub>3</sub>): δ 9.1 (s, 1H, Thymine aromatic), 7.7-7.2 (m, 8H, Florenyl aromatic protons), 6.7 (s, 1H, Thymine-NH-), 5.9 (br.s, 1H, fmoc-HN-), 5.8 (m, 1H, -CH=CH<sub>2</sub>), 5.3 (s, 1H), 5.25 (s, 2H), 5.2 (m, 2H), 5 (m, 2H), 4.6 (m, 2H), 4.4 (m, 2H), 4.2-4 (m, 2H), 3.8 (d, 1H, J=6.4 Hz), 3.7 (q, 1H, J=6.4 Hz), 3.4 (m, 2H), 3 (m, 1H), 2.9 (m, 1H), 2.2-2 (3s, 9H, sugar acetyl protons), 1.9 (s, 3H, Thymine-CH<sub>3</sub>), 1.1 (s, 3H, fucose-CH<sub>3</sub>).

15 HRMS (M+H)<sup>+</sup>, calcd (found) for C<sub>44</sub>H<sub>50</sub>N<sub>5</sub>O<sub>15</sub> is 876.3303(876.3304).

#### EXAMPLE 34

##### **Fmoc-N-(Thymine-1-ylacetyl)Asp-[O-(2,3,4,5-tetra-O-acetyl glucosamine-1-yl)-amide]-O-Allyl (36)**

20 *Preparation: procedure (e).* Yield = 73%.

Mp = ;  $[\alpha]_D^{22}$  (c=0.5, methanol); MS (FAB) *m/z* 933 (M+H); <sup>1</sup>H-NMR, (CDCl<sub>3</sub>): δ 10 (s, 1H, Thymine aromatic), 7.7-7.2 (m, 8H, Florenyl aromatic protons), 6.8 (s, 1H, Thymine-NH-), 6 (br.s, 1H, fmoc-HN-), 5.8 (m, 1H, -CH=CH<sub>2</sub>), 5.4-5 (m, 5H), 4.7 (m, 1H), 4.6 (m, 2H), 4.4 (m, 2H), 4.2 (m, 3H), 4.1 (m, 3H), 3.8-3.2 (m, 6H), 2.1-2 (4s, 12H, sugar acetyl protons), 1.9 (s, 3H, Thymine-CH<sub>3</sub>); <sup>13</sup>C-NMR (CDCl<sub>3</sub>): δ 171.78, 171.08, 170.97, 170.71, 170.51, 169.42, 169.23, 166.94, 164.52, 156.47, 151.12, 143.58, 141.06, 131.05, 130.79 (-CH=CH<sub>2</sub>), 127.60, 126.92, 124.82, 119.84, 119.51, 119.17 (-CH=CH<sub>2</sub>), 110.40, 79.39, 73.07, 67.89, 66.48, 61.68, 57.94, 52.75, 48.96, 48.28, 47.00, 39.14, 35.51, 33.67, 24.73, 22.74, 20.54, 20.43, (sugar-CO-CH<sub>3</sub> groups), 12.11 (Thymine-CH<sub>3</sub>).

25 C.H.N analyse for C<sub>45</sub>H<sub>52</sub>N<sub>6</sub>O<sub>16</sub>, 3/2 H<sub>2</sub>O: calc. C 56.30, H 5.78, N 8.75; found C 56.43, H 5.70, N 8.72.

#### EXAMPLE 35

##### **Fmoc-N-(Thymine-1-ylacetyl)Asp-[O-(2,3,4,5-tetra-O-acetyl galactosamine-1-yl)-amide]-O-Allyl (37)**

35

*Preparation: procedure (e).* Yield = 60%.

Mp = ;  $[\alpha]_D^{22}$  (c=0.5, methanol); MS (FAB)  $m/z$  933 (M+H);  $^1\text{H-NMR}$ , ( $\text{CDCl}_3$ ):  $\delta$  10.2 (s, 1H, Thymine aromatic), 7.9-7 (m, 8H, Florenyl aromatic protons), 6.8 (s, 1H, Thymine-NH-), 6.3 (m, 1H), 6.1 (br.s, 1H, fmoc-HN-), 5.8 (m, 1H,  $-\text{CH}=\text{CH}_2$ ), 5.4-5.1 (m, 4H), 4.7 (m, 2H), 4.4 (m, 2H), 4.2 (m, 5H), 3.6-3.3 (m, 4H), 2.8 (m, 4H), 2.1-2 (m, 12H, sugar acetyl protons), 1.8 (s, 3H, Thymine-CH<sub>3</sub>);  $^{13}\text{C-NMR}$  ( $\text{CDCl}_3$ ):  $\delta$  171.78, 171.08, 170.97, 170.71, 170.51, 169.42, 169.23,  $^{13}\text{C-NMR}$  ( $\text{CDCl}_3$ ):  $\delta$  172.32, 170.82, 170.54, 170.33, 170.04, 169.55, 169.37, 167.03, 164.53, 156.62, 151.35, 143.55, 141.04, 131.09, 127.59, 126.91, 124.80, 119.82, 119.07( $-\text{CH}=\text{CH}_2$ ), 110.41, 79.57, 71.83, 70.46, 67.91, 67.37, 66.86, 66.70, 66.45, 61.77, 61.31, 58.59, 49.20, 48.98, 47.96, 47.01, 39.18, 23.04, 22.95, 22.77, 20.52 (sugar-CO-CH<sub>3</sub> groups), 12.04 (Thymine-CH<sub>3</sub>).

HRMS (M+H)<sup>+</sup>, calcd (found) for C<sub>45</sub>H<sub>53</sub>N<sub>6</sub>O<sub>16</sub> is 933.3518(933.3549).

#### EXAMPLE 36

##### 4-C-(2,3,4,6-tera-O-acetyl-galactose-1-yl)-2-Z-amino-2-butenic acid t-butyl ester (41)

*Preparation:* 2.2 g (5.8 mmol, 1.1 eq) **40** was dissolved in 20 ml dry THF. 0.8 ml (mmol, 1.1 eq) tetramethylguanidine was added at (−78°C) and stirred for 5 min. 1.9 g (5.27 mmol, 1 eq) 2,3,4,5 tetraacetyl galactose-1yl-acetaldehyde was added in 10 ml dry THF solution. The reaction mixture was allowed to reach the room temperature. Volatile was evaporated and the residue was purified on silica gel column eluting hexane-ethylacetate 1:1 solution. 2.2 g (3.5 mmol) titled compound was obtained as colourless oil. Yield = 66%.

Mp: ;  $[\alpha]_D^{22}$  (c=0.5, methanol); MS (FAB)  $m/z$  622 (M+H);  $^1\text{H-NMR}$ , ( $\text{CDCl}_3$ ):  $\delta$  7.4 (m, 4H of cbz aromatic protons), 6.5 (overlapping, 2H, cbz-ring-<sup>4</sup>CH-, cbz-HN-), 5.4 (t, 1H, J=3 Hz), 5.2 (dd, 1H, J=9.1, 4.8 Hz), 5.18 (dd, 1H, J=9.1, 3.3 Hz), 5.15 (d, 2H, J=1.3 Hz), 4.35 (m, 1H), 4.45 (m, 1H), 4.1-4 (m, 2H), 2.6 (m, 1H), 2.4 (m, 1H), 2.18 (d, 1H, J=8.4), 2.15-2 (4s, 12H, sugar acetyl protons), 1.4 (s, 9H, t-Bu);  $^{13}\text{C-NMR}$  ( $\text{CDCl}_3$ ):  $\delta$  170.42, 169.86, 169.68, 163.08, 153.86, 135.81, 129.60, 128.43, 128.16, 128.06, 82.06, 74.04, 71.02, 68.91, 68.51, 67.72, 67.55, 67.32, 67.25, 61.18, 27.87, 25.82, 20.62, 20.54.

HRMS (M+H)<sup>+</sup>, calcd (found) for C<sub>30</sub>H<sub>40</sub>NO<sub>13</sub> is 622.2500(622.2495).

#### EXAMPLE 37

##### 4-C-(2,3,4,6-tera-O-acetyl-galactose-1-yl)-2(2-fmoc-aminoethyl)-aminobutanoic acid t-butyl ester (43)

*Preparation:* 2.2 g (3.5 mmol) **41** was dissolved in 10 ml methanole. 50 mg Pd/C catalyst was added and the mixture hydrogenated at 1 atm for 3 hours. The catalyst was filtered off over celite and methanol was removed under vacuum. The remaining oil **42** was directly used for procedure (b). 2.1g (2.78 mmol) titled compound was obtained as white solid. Yield = 82%.

Mp = ;  $[\alpha]_D^{22} = (c=0.5, \text{methanol})$ ; MS (FAB)  $m/z$  755 (M+H);  $^1\text{H-NMR}$ , ( $\text{CDCl}_3$ ):  $\delta$  7.7-7.2 (m, 8H, fmoc aromatic protons), 5.5 (overlapping s, 2H), 5.2 (m, 2H), 4.4 (d, 2H, 7 Hz), 4.2 (s, 2H), 4.1 (m, 2H), 3.3-3 (m, 4H), 2.8 (m, 1H), 2.6 (m, 1H), 2.2-2 (4s, 12H, sugar acetyl protons), 1.4(s, 9H, t-Bu);  $\text{CDCl}_3$   $^{13}\text{C-NMR}$  ( $\text{CDCl}_3$ ):  $\delta$  174.13, 170.53, 170.47, 170.40, 170.01, 169.93, 169.86, 169.76, 169.71, 169.63, 156.37, 143.79, 143.69, 141.07, 127.45, 126.82, 124.90, 119.75, 81.43, 71.60, 68.26, 68.20, 68.00, 67.80, 67.71, 67.58, 67.44, 67.35, 66.51, 61.42, 61.24, 61.07, 61.00, 60.19, 47.33, 47.05, 28.79, 27.89, 27.47, 21.75, 20.86, 20.59, 20.53, 20.46 (sugar-CO-CH<sub>3</sub> groups), 14.01.

HRMS (M+H)<sup>+</sup>, calcd (found) for C<sub>39</sub>H<sub>51</sub>N<sub>2</sub>O<sub>13</sub> is 755.3391 (755.3371).

#### EXAMPLE 38

##### **4-C-(2,3,4,6-tera-O-acetyl-galactose-1-yl)-2[N,N(2-fmoc-aminoethyl)-(thymine-1-yl-methyl-carbonyl)-aminobutanoic acid t-butyl ester (44)**

*Preparation: procedure (c).* Yield = 66%.

Mp: ;  $[\alpha]_D^{22} = (c=0.5, \text{methanol})$ ; MS (FAB)  $m/z$  951 (M+H);  $^1\text{H-NMR}$ , ( $\text{CDCl}_3$ ):  $\delta$  9.2 (s, 1H, thymine aromatic), 7.8-7.3 (m, 8H, fmoc aromatic), 6.9 (s, 1H, thymine-NH-), 5.8 (s, 1H, fmoc-HN-), 5.4 (s, 1H), 5.35 (dd, 1H, J=10.2, 3.3 Hz), 5.25 (dd, 1H, J=10.2, 5.7 Hz), 5.2 (d, 1H, J=3 Hz), 4.4 (m, 3H), 4.2-4 (m, 5H), 3.6-3.3 (m, 4H), 2.2-2 (4s, 12H, sugar acetyl protons), 1.8 (s, 3H, thymine-methyl), 1.4 (s, 9H, t-Bu);  $^{13}\text{C-NMR}$  ( $\text{CDCl}_3$ ):  $\delta$  170.46, 169.94, 169.84, 169.07, 168.19, 166.85, 163.99, 156.44, 150.82, 143.54, 141.08, 127.59, 126.89, 124.78, 119.83, 110.23, 82.49, 72.31, 71.83, 67.49, 66.93, 66.61, 61.81, 61.29, 60.20, 59.30, 48.87, 48.40, 47.89, 46.96, 46.10, 39.98, 39.38, 33.72, 27.75, 25.41, 24.88, 22.60, 21.16, 20.86, 20.68, 20.55, 14.01, 12.16.

HRMS (M+H)<sup>+</sup>, calcd (found) for C<sub>46</sub>H<sub>57</sub>N<sub>4</sub>O<sub>18</sub> is 921.3770(921.3782).

#### EXAMPLE 39

##### **N-Fmoc-N-(Thymine-1-ylacetyl)Serine(2,3,4,5-tetra-O-acetyl- $\alpha$ -D-Galactose-1-yl)-COOH (1s)**

*Preparation: procedure (f).* Yield = 86%.

Mp = 124-127;  $[\alpha]_D^{22} = (c=0.5, \text{methanol})$ ; MS (FAB)  $m/z$  867 (M+H);  $^1\text{H-NMR}$ , ( $\text{CD}_3\text{OD}$ ): characteristic signals:  $\delta$  = 7.8-7.3 [m, 9H, (8H, Florenyl aromatic protons, 1H, Thymine aro-

matic)], 5.4 (s, 1H, sugar <sup>4</sup>CH), 5.15 (m, 2H, sugar <sup>3</sup>CH and <sup>2</sup>CH), 4.65 (overlapping s, 3H, sugar <sup>1</sup>CH, -CO-CH<sub>2</sub>-Thymine), 2.2-2 (4s, 12H, sugar acetyl protons), 1.8 (s, 3H, Thymine-CH<sub>3</sub>); <sup>13</sup>C-NMR (CD<sub>3</sub>OD): δ 170.20, 170.13, 170.05, 169.75, 165.15, 143.56, 143.46, 141.98, 140.73, 133.95, 128.16, 127.99, 126.92, 126.36, 126.30, 124.40, 124.30, 119.06, 108.83, 100.26 (sugar C1), 70.33, 70.13, 68.53, 68.44, 66.90, 65.86, 60.58, 47.95, 38.61, 18.88, 18.68, 18.59, 10.36.

HRMS (M+H)<sup>+</sup>, calcd (found) for C<sub>41</sub>H<sub>47</sub>N<sub>4</sub>O<sub>17</sub> is 867.2936 (867.2924).

#### EXAMPLE 40

##### 10 ***N*-Fmoc-*N*-(Thymine-1-ylacetyl)Threonine(2,3,4,5-tetra-*O*-acetyl-α-D-Galactose-1-yl)-COOH (2s)**

*Preparation: procedure (f).* Yield = 90%.

Mp = 128-131; [α]<sub>D</sub><sup>22</sup> = (c=0.5, methanol); MS (FAB) *m/z* 881 (M+H); <sup>1</sup>H-NMR, (CD<sub>3</sub>OD): characteristic signals: δ = 7.84-7.32 [m, 9H, (8H, Florenyl aromatic protons, 1H, Thymine aromatic), 5.41 (d, 1H, J=3.3 Hz, sugar <sup>4</sup>CH), 5.2 (dd, 1H, J=10.44, 3.3 Hz, sugar <sup>3</sup>CH), 5.1 (dd, 1H, J<sub>1,2</sub>=7.5, J<sub>2,3</sub>=10.5, sugar <sup>2</sup>CH), 2.18-2 (4s, 12H, sugar acetyl protons), 1.9 (s, 3H, Thymine-CH<sub>3</sub>), 1.36 (d, 3H, J=5.86 Hz, <sup>β</sup>C-CH<sub>3</sub>); <sup>13</sup>C-NMR (CD<sub>3</sub>OD): δ = 172.06, 171.51, 167.09, 145.38, 143.86, 142.61, 128.87, 128.26, 126.26, 121.03, 116.69, 110.78, 100.40(sugar C1), 74.76, 72.31, 71.97, 70.66, 68.80, 67.78, 62.69, 62.49, 20.93, 20.73, 20.62, 12.41.

HRMS (M+H)<sup>+</sup>, calcd (found) for C<sub>42</sub>H<sub>49</sub>N<sub>4</sub>O<sub>17</sub> is 881.3043 (881.3073).

#### EXAMPLE 41

##### 25 ***N*-boc-*N*-(Thymine-1-ylacetyl)Lisine-[*N*-*N*-bis(2,3,4,5-tetra-*O*-acetyl Galactose-1-yl)-ethyl]-OH (3s)**

*Preparation: procedure (f).* Yield = 80%.

Mp = 118-120; [α]<sub>D</sub><sup>22</sup> = (c=0.5, methanol); MS (FAB) *m/z* 1172 (M+H); <sup>1</sup>H-NMR, (CD<sub>3</sub>OH): resolved signals: δ 7.4 (s, 1H, aromatic proton of Thymine), 5.45 (d, 2H, J=2.74 Hz, 2 x <sup>4</sup>CH-sugar), 5.31 (dd, 2H, J=9.16, 3.3 Hz, 2 x <sup>3</sup>CH-sugar), 5.25 (m, 2H, 2 x <sup>2</sup>CH-sugar), 4.38 (m, 4H), 4.25 (m, 2H), 4.15 (m, 3H), 3.95 (dd, 1H, J=12.82, 3.66Hz), 3.65 (dt, 1H, J=2.38, 12.45 Hz), 3.5 (m, 2H), 3.15 (d, 2H, J=13.7 Hz), 2.9 (br, 2H), 2.8(m, 2H), 2.2-2 (4s, 24H, CO-CH<sub>3</sub> of sugar), 1.9 (s, 3H, CH<sub>3</sub> of Thymine), 1.8-1.6 (br. 4H), 1.47 (overlapping s, 11H, 9H of boc, 2H of lysine side chain); <sup>13</sup>C-NMR (CD<sub>3</sub>OH): δ 172.41, 172.23, 171.91, 171.72, 171.69, 171.50, 170.39, 169.75, 167.08, 158.28, 153.16, 144.45, 144.16, 110.77, 110.57, 80.42, 71.31, 71.24, 70.29, 69.86, 69.78, 69.37, 69.20, 69.07, 66.30, 65.76, 62.88, 62.70, 54.96, 51.67,

50.87, 45.10, 40.60, 39.83, 32.81, 31.39, 30.56, 28.95, 28.88, 26.81, 26.26, 25.95, 25.66, 23.46, 23.01, 20.99, 20.79, 20.68, 12.43.

HRMS (M+H)<sup>+</sup>, calcd (found) for C<sub>52</sub>H<sub>73</sub>N<sub>5</sub>O<sub>26</sub> is 1172.4986 (1172.5027).

## 5 EXAMPLE 42

***N*-boc-*N*-(Thymine-1-ylacetyl)Lisine-[*N*-*N*-bis(2,3,4-tri-*O*-acetyl fucose-1-yl)-ethyl]-OH (4s)**

*Preparation: procedure (f).* Yield = 88%.

Mp = ; [α]<sub>D</sub><sup>22</sup> = (c=0.5, methanol); MS (FAB) *m/z* 1056 (M+H); <sup>13</sup>C-NMR (CD<sub>3</sub>OH): δ 172.19, 171.65, 166.98, 162.58, 158.30, 153.14, 144.9, 133.86, 133.18, 133.05, 130.66, 130.13, 128.83, 128.35, 123.36, 114.36, 110.83, 108.98, 80.44, 80.08, 71.82, 71.51, 69.82, 69.34, 67.49, 62.40, 54.33, 51.85, 51.32, 50.66, 45.12, 40.60, 39.81, 30.22, 28.97, 28.90, 25.24, 22.13, 21.02, 20.83, 20.67, 16.43, 12.49, 11.99.

HRMS (M+H)<sup>+</sup>, calcd (found) for C<sub>46</sub>H<sub>74</sub>N<sub>5</sub>O<sub>21</sub> is 1056.4876(1056.4886).

***N*-boc-*N*-(Thymine-1-ylacetyl)Lisine-[*N*-(2,3,4,5-tetra-*O*-acetyl Galactose-1-yl)-acetyl]-OH (5s)**

*Preparation: procedure (f).* Yield = 95%.

Mp = 115-118; [α]<sub>D</sub><sup>22</sup> = (c=0.5, methanol); MS (FAB) *m/z* 828 (M+H); <sup>1</sup>H-NMR, (CD<sub>3</sub>OD): resolved signals: δ 7.37 (s, 1H, aromatic proton of Thymine), 5.44 (d, 1H, J=2.56 Hz, <sup>4</sup>CH-sugar), 5.3 (overlapping, 2H, <sup>3</sup>CH- and <sup>2</sup>CH-sugar), 4.7 (m, 2H), 4.5(m, 1H), 4.25 (m, 1H), 4.1 (m, 3H), 3.5 (m, 1H), 3.25 (m, 2H), 2.7 (m, 1H), 2.5 (dd, 1H, J=5.3, 2.2 Hz), 2.2-2 (4s, 12H, CO-CH<sub>3</sub> of sugar), 1.9 (s, 3H, CH<sub>3</sub> of Thymine), 1.6 (br. 2H), 1.4 (overlapping s, 11H, 9H of boc, 2H of lysine side chain); <sup>13</sup>C-NMR (CD<sub>3</sub>OD): δ 173.13, 172.16, 170.41-169.52 (sugar-O-CO-Me), 167.69, 164.97 (-CO-CH<sub>2</sub>-Thymine), 156.48 (-NH-COO-t-butyl), 151.46, 141.90, 109.37, 78.68 [-O-C(Me)<sub>3</sub>], 74.39 (sugar C5), 69.32-66.93 (sugar C1-4), 60.52 (sugar C6), 48.68 (lysine, α carbon), 40.93, 38.31, 37.08, 33.06, 29.22-25.82 [-COO-C(CH<sub>3</sub>)<sub>3</sub>], 22.94-18.65 (sugar-O-CO-CH<sub>3</sub>), 10.34 (Thymine-CH<sub>3</sub>).

HRMS (M+H)<sup>+</sup>, calcd (found) for C<sub>36</sub>H<sub>54</sub>N<sub>5</sub>O<sub>17</sub> is 828.3515 (828.3515).

## EXAMPLE 43

***N*-fmoc-*N*-(Thymine-1-ylacetyl)Lisine-[*N*-(2,3,4,5-tetra-*O*-acetyl Galactose-1-yl)-acetyl]-OH (6s)**

*Preparation: procedure (f).* Yield = 83%.



Mp = 130-132;  $[\alpha]_D^{22}$  (c=0.5, methanol); MS (FAB)  $m/z$  951 (M+H);  $^1\text{H-NMR}$ , ( $\text{CD}_3\text{OH}$ ): resolved signals:  $\delta$  7.8-7.3 (9H, aromatic proton of Thymine, 8H of florenyl aromatics), 5.43 (d, 1H,  $J=2.4$  Hz,  $^4\text{CH-sugar}$ ), 5.3 (overlapping, 2H,  $^3\text{CH-}$  and  $^2\text{CH-sugar}$ ), 4.72 (m, 3H), 4.45 (d, 1H,  $J=6.6$ ), 4.4 (m, 2H), 4.3-4.2 (m, 3H), 4.2-4.1 (m, 3H), 3.6 (m, 1H), 3.45 (m, 2H), 3.2 (m, 3H), 2.7 (dd, 1H,  $J=14.65$ , 9.34 Hz), 2.5 (dd, 1H,  $J=14.65$ , 5.31Hz), 2.2-2 (3s, 12H, CO-CH<sub>3</sub> of sugar), 1.9 (s, 3H, CH<sub>3</sub> of Thymine), 1.6 (br. 3H), 1.2 (m, 1H);  $^{13}\text{C-NMR}$  ( $\text{CD}_3\text{OH}$ ):  $\delta$  174.33, 172.23, 172.20, 171.89, 171.56, 171.41, 170.43, 169.78, 166.98, 158.89, 153.02, 145.02, 143.91, 142.67, 128.88, 128.23, 126.22, 121.04, 110.92, 71.28, 70.14, 69.41, 69.15, 69.00, 67.81, 62.52, 61.65, 50.20, 40.80, 40.30, 35.08, 30.12, 29.78, 24.96, 20.83, 20.76, 20.72, 20.62, 12.37(Thymine-CH<sub>3</sub>).

HRMS (M+H)<sup>+</sup>, calcd (found) for C<sub>46</sub>H<sub>56</sub>N<sub>5</sub>O<sub>17</sub> is 950.3671 (950.3666).

#### EXAMPLE 44

***N*-fmoc-*N*-(Thymine-1-ylacetyl)Lysine-[*N*-(2,3,4,5-tetra-*O*-acetyl mannose-1-yl)-acetyl]-*O*H (7s)**

*Preparation: procedure (f). Yield = 92%.*

Mp = ;  $[\alpha]_D^{22}$  (c=0.5, methanol); MS (FAB)  $m/z$  951 (M+H);  $^{13}\text{C-NMR}$  ( $\text{CDCl}_3$ ):  $\delta$  193.55, 178.08, 172.71, 172.46, 171.75, 171.56, 171.44, 166.88, 153.09, 145.27, 144.01, 142.58, 130.33, 128.89, 128.24, 126.23, 121.04, 116.26, 110.95, 73.18, 72.40, 72.30, 71.40, 70.19, 68.28, 67.88, 63.44, 62.41, 40.15, 37.31, 30.17, 25.05, 20.87, 20.71(acetyl groups on sugar), 12.40 (CH<sub>3</sub> of Thymine)

#### EXAMPLE 45

***N*-fmoc-*N*-(Thymine-1-ylacetyl)Lysine-[*N*-(2,3,4-ttri-*O*-acetyl fucose-1-yl)-acetyl]-*O*H (8s)**

*Preparation: procedure (f). Yield = 86%.*

Mp = ;  $[\alpha]_D^{22}$  (c=0.5, methanol); MS (FAB)  $m/z$  892 (M+H);  $^1\text{H-NMR}$ , ( $\text{CD}_3\text{OD}$ ): resolved protons  $\delta$  7.8-7.2 (m, 8H, Florenyl aromatic protons), 6.9 (s, 1H, Thymine-NH-), 5.3 (overlapping s, 2H), 4.7 (s, 2H, -CO-CH<sub>2</sub>-Thymine), 4.44 (d, 1H,  $J=6.6\text{Hz}$ ), 4.38 (m, 1H), 4.25 (m, 1H), 4.15 (m, 1H), 3.6-3.2 (multiplets, 5H), 2.7 (m, 1H), 2.5 (m, 1H), 2.2-2 (3s, 9H, sugar acetyl protons), 1.85 (s, 3H, Thymine-CH<sub>3</sub>), 1.6-1.3 (m, 4H, lysine side chain protons), 1.1 (d, 3H,  $J=4.2$ , fucose-CH<sub>3</sub>);  $^{13}\text{C-NMR}$  ( $\text{CD}_3\text{OD}$ ):  $\delta$  174.67, 172.51, 172.24, 171.51, 170.32, 169.68, 166.93, 158.86, 152.99, 145.29, 143.93, 142.62, 133.19, 131.77, 130.48, 129.92, 128.89, 128.24, 126.22, 121.83, 121.05, 117.01, 110.89, 72.044, 71.53, 69.88, 69.00, 67.81,

61.93, 42.08, 40.75, 40.12, 34.79, 30.15, 29.75, 26.77, 26.09, 24.91, 20.80, 20.59, 16.50, 12.38.

HRMS (M+H)<sup>+</sup>, calcd (found) for C<sub>44</sub>H<sub>54</sub>N<sub>5</sub>O<sub>15</sub> is 892.3616(892.3635).

#### 5 EXAMPLE 46

**fmoc-N-(Thymine-1-ylacetyl)Asp-[O-(2,3,4,5-tetra-O-acetyl Galactose-1-yl)-amide]-OH (9s)**

*Preparation: procedure (f).* Yield = 94%.

Mp = ; [α]<sub>D</sub><sup>22</sup> = (c=0.5, methanol); MS (FAB) *m/z* 894 (M+H); <sup>1</sup>H-NMR, (CD<sub>3</sub>OD): resolved protons δ 7.8-7.3 (m, 8H, Florenyl aromatic protons), 7.2 (s, 1H, Thymine-NH-), 5.45 (s, 1H, sugar <sup>4</sup>CH), 5.37 (d, 1H, J=8.5 Hz, <sup>2</sup>CH), 4.24(m, 2H), 5.23 (d, 1H, J=13.4 Hz), 4.48 (t, 1H, J=6 Hz), 4.42 (d, 1H, J=3.3 Hz, -CO-CH<sub>2</sub>-Thymine), 4.3-4.1 (m, 3H), 2.2-2 (4s, 12H, sugar acetyl protons), 1.8 (s, 3H, Thymine-CH<sub>3</sub>); <sup>13</sup>C-NMR (CD<sub>3</sub>OD): δ 173.62, 172.93, 172.00, 171.79, 171.53, 169.34, 166.98, 158.89, 152.90, 145.33, 143.68, 142.64, 133.85, 133.21, 130.14, 128.89, 128.27, 126.29, 121.06, 110.93, 79.30, 73.41, 73.09, 69.81, 69.03, 68.36, 62.72, 59.72, 40.46, 37.15, 34.80, 32.77, 30.19, 26.79, 26.10, 24.91, 23.73, 20.94, 20.70, 20.64, 14.51, 12.38.

HRMS (M+H)<sup>+</sup>, calcd (found) for C<sub>42</sub>H<sub>48</sub>N<sub>5</sub>O<sub>17</sub> is 894.3045 (894.3072).

#### 20 EXAMPLE 47

**Fmoc-N-(Thymine-1-ylacetyl)Asp-[O-(2,3,4,5-tetra-O-acetyl mannose-1-yl)-amide]-OH (10s)**

*Preparation: procedure (f).* Yield = 95%.

Mp = ; [α]<sub>D</sub><sup>22</sup> = (c=0.5, methanol); MS (FAB) *m/z* 894 (M+H); <sup>1</sup>H-NMR, (DMSO): resolved protons δ 11.2 (s, 1H, thymine aromatic), 8.7 (d, 1H, J=8.8 Hz, Thymine-NH-) 7.8-7.3 (m, 8H, Florenyl aromatic protons), 5.6 (d, 1H, J=8.8 Hz), 5.37 (dd, 1H, J=10.2, 3.8 Hz), 5.15 (d, 1H, J=3.3 Hz), 5 (t, 1H, J=10 Hz), 4.6 (m, 2H), 4.3 (d, 1H, J=6 Hz), 4.2 (m, 2H), 4 (d, 1H, J=11 Hz, 2.2-2 (4s, 12H, sugar acetyl protons), 1.8 (s, 3H, Thymine-CH<sub>3</sub>); (CDCl<sub>3</sub>)<sup>13</sup>C-NMR (CDCl<sub>3</sub>): δ 170.13, 169.62, 143.91, 140.76, 127.67, 127.13, 125.17, 120.15, 108.05, 72.60, 70.88, 65.58, 47.84, 46.75, 33.39, 24.53, 21.03, 20.61, (sugar-CO-CH<sub>3</sub> groups), 11.99 (Thymine-CH<sub>3</sub>).

HRMS (M+H)<sup>+</sup>, calcd (found) for C<sub>42</sub>H<sub>48</sub>N<sub>5</sub>O<sub>17</sub> is 894.3045 (893.3065).

#### EXAMPLE 48

35 **Fmoc-N-(Thymine-1-ylacetyl)Asp-[O-(2,3,4-tri-O-acetyl fucose-1-yl)-amide]-OH (11s)**

*Preparation: procedure (f).* Yield = 88%.

Mp = ;  $[\alpha]_D^{22}$  (c=0.5, methanol); MS (FAB)  $m/z$  (M+H); 7.8-7.3 (m, 8H, Florenyl aromatic protons), 7.2 (s, 1H, Thymine-NH-), 5.3 (overlapping s, 2H), 5.2 (d, 1H, J=3 Hz), 4.66 (s, 1H), 4.4 (m, 2H), 4.25 (m, 1H), 4 (m, 1H), 3.5 (m, 3H), 3.15 (m, 2H), 2.2-2 (singles, 9H, sugar acetyl protons), 1.8 (s, 3H, Thymine-CH<sub>3</sub>), 1.1 (s, 3H, fucose-CH<sub>3</sub>); <sup>13</sup>C-NMR (CD<sub>3</sub>OD):  $\delta$  172.30, 171.81, 170.68, 170.30, 169.96, 167.74, 165.39, 157.35, 151.41, 143.84, 143.74, 142.15, 141.04, 134.35, 132.27, 131.60, 131.49, 128.52, 128.40, 127.30, 126.68, 124.77, 124.07, 119.44, 109.36, 86.67, 77.61, 71.92, 71.46, 71.33, 70.51, 68.57, 68.26, 68.12, 67.93, 66.40, 58.96, 39.00, 35.92, 33.26, 25.26, 24.57, 19.27, 19.11, 19.05, 15.05, 10.90.

HRMS (M+H)<sup>+</sup>, calcd (found) for C<sub>40</sub>H<sub>46</sub>N<sub>5</sub>O<sub>15</sub> is 836.2990 (836.2980).

#### EXAMPLE 49

##### **Fmoc-N-(Thymine-1-ylacetyl)Asp-[O-(2,3,4,5-tetra-acetyl glucosamine-1-yl)-amide]-OH (12s)**

*Preparation: procedure (f).* Yield = 90%.

Mp = ;  $[\alpha]_D^{22}$  (c=0.5, methanol); MS (FAB)  $m/z$  (M+H); <sup>1</sup>H-NMR, (CD<sub>3</sub>OD):  $\delta$  7.8-7.3 (m, 8H, Florenyl aromatic protons), 7.2 (d, 1H, J=9Hz, Thymine-NH-), 5.3 (m, 2H), 5.1 (d, 1H, J=9.7Hz), 4.7 (m, 1H), 4.4 (m, 2H), 4.2 (m, 2H), 4.1 (m, 1H), 3.9 (m, 1H), 3.6-2.8 (multiplets, 5H), 2.1-1.9 (4s, 12H, sugar acetyl protons), 1.8 (s, 3H, Thymine-CH<sub>3</sub>); <sup>13</sup>C-NMR (CD<sub>3</sub>OD):  $\delta$  173.76, 172.94, 172.43, 172.05, 171.40, 170.14, 169.26, 166.94, 158.85, 153.13, 152.87, 145.32, 143.72, 142.59, 130.13, 129.93, 128.88, 128.26, 126.24, 121.05, 111.17, 110.92, 79.53, 74.85, 74.62, 69.97, 69.82, 67.86, 63.34, 59.82, 54.28, 40.46, 37.16, 23.09, 23.01, 22.87, 22.74, 20.72, 12.39.

HRMS (M+H)<sup>+</sup>, calcd (found) for C<sub>42</sub>H<sub>48</sub>N<sub>5</sub>O<sub>18</sub> is 893.3205 (893.3232).

#### EXAMPLE 50

##### **Fmoc-N-(Thymine-1-ylacetyl)Asp-[O-(2,3,4,5-tetra-acetyl galactosamine-1-yl)-amide]-OH (13s)**

*Preparation: procedure (f).* Yield = 91%.

Mp = ;  $[\alpha]_D^{22}$  (c=0.5, methanol); MS (FAB)  $m/z$  (M+H); <sup>1</sup>H-NMR, (CD<sub>3</sub>OD):  $\delta$  7.8-7.3 (m, 8H, Florenyl aromatic protons), 7.2 (s, 1H, Thymine-NH-), 5.4 (s, 1H), 5.3 (d, 1H, J=9.5 Hz), 5.2 (dd, 1H, J=11, 3.3 Hz), 5.1 (m, 1H), 4.7 (s, 1H), 4.4 (m, 2H), 4.3-4.1 (m, 4H), 3.8 (m, 1H), 3.6 (m, 1H), 3.5-2.8 (multiplets, 5H), 2.2-1.9 (4s, 12H, sugar acetyl protons), 1.85 (s, 3H, Thymine-CH<sub>3</sub>); <sup>13</sup>C-NMR (CD<sub>3</sub>OD):  $\delta$  174.06, 173.57, 172.94, 172.18, 172.05, 171.81, 171.74, 169.28, 166.98, 158.88, 153.14, 152.9, 145.34, 143.75, 142.62, 135.10, 134.97,

133.05, 131.43, 130.22, 129.95, 128.87, 128.24, 126.24, 121.03, 110.89, 80.06, 73.36, 72.71, 68.91, 68.23, 67.89, 62.86, 59.86, 40.46, 37.19, 26.55, 23.12, 20.70, 12.39.

HRMS (M+H)<sup>+</sup>, calcd (found) for C<sub>42</sub>H<sub>49</sub>N<sub>6</sub>O<sub>16</sub> is 893.3205 (893.3245).

## 5 EXAMPLE 51

### **4-C-(2,3,4,6-tera-O-acetyl-galactose-1-yl)-2[N,N(2-fmoc-aminoethyl)-(thymine-1-yl-methyl-carbonyl)-aminobutanoic acid (14s)**

920 mg (1 mmol) **44** was added to 10 ml solution of 5% triethylsilane in trifluoroacetic acid at 0°C and stirred until TLC did not show any starting material left. 50 ml DCM was added and volatile removed under vacuum. Further 3 x 100 ml DCM was added and evaporated in order to removal of TFA. The product was precipitated with adding diethylether. 830 mg (0.94 mmol) titled compound was obtained as white powder (Yield = 94%).


Mp= ; [α]<sub>D</sub><sup>22</sup>= (c=0.5, methanol); MS (FAB) *m/z* 865 (M+H); <sup>1</sup>H-NMR, (CD<sub>3</sub>OH): δ 7.8-7.3 (m, 8H, fmoc aromatic), 7.2 (s, 1H, thymine-NH-), 5.4 (overlapping s, 2H), 5.3 (m, 1H), 4.7(s, 1H), 4.6 (m, 1H), 4.5(d, 2H, J=6.6 Hz), 4.2 (overlapping s, 2H), 4.1 (overlapping s, 2H), 3.6 (s, 1H), 3.5 (overlapping s, 2H), 3.3 (s, 1H), 2.2-2 (4s, 12H, sugar acetyl protons), 1.85 (s, 3H, thymine-methyl), <sup>13</sup>C-NMR (CD<sub>3</sub>OD): δ 172.75, 171.22, 170.86, 170.75, 170.63, 170.52, 168.56, 165.80, 157.63, 151.63, 144.09, 142.89, 142.70, 141.43, 127.66, 127.01, 125.00, 119.82, 109.43, 72.16, 68.62, 68.45, 68.26, 68.08, 39.52, 33.60, 31.59, 25.58, 24.91, 24.76, 22.55, 21.73, 19.60, 19.43, 13.31, 11.16.

HRMS (M+H)<sup>+</sup>, calcd (found) for C<sub>42</sub>H<sub>49</sub>N<sub>4</sub>O<sub>16</sub> is 865.3140 (865.3144).

## EXAMPLE 52

### **Solid Phase Synthesis of Type IV PNA monomers**

52



#### *N<sup>α</sup>-Boc-Diaminopropionic acid*

- 5 A) To a solution of Boc-L-Asn-OH (15.0 g, 60.3 mmol) in EtOAc (75 mL), CH<sub>3</sub>CN (75 mL) and water (30 mL) at 10°C was added PIDA (24.9 g, 77.3 mmol, 1.3 eq.). The mixture was stirred for 30 min, heated to 20°C and stirred for further 2.5 h. The mixture was heated to 70°C until completely dissolved and then slowly cooled to 20°C and filtered. The remanence was washed with EtOAc (2 x 50 mL) and dried. Yield 8.9 g (68%). <sup>1</sup>H NMR (DMSO-*d*<sub>6</sub> + TFA)
- 10 δ 8.06 (bs, 3NH), 7.36 (d, 1NH, *J* = 8.7 Hz), 4.40–4.25 (m, 1H), 3.35–3.25 (m, 1H), 3.15–3.00 (m, 1H), 1.51 (s, 9H). MS: [M+H]<sup>+</sup>: expected: 205.1; observed: 205.1.

#### *N<sup>α</sup>-Boc-N<sup>β</sup>-Z-Diaminopropionic acid*

- 15 B) To a solution of *N<sup>α</sup>-Boc-Diaminopropionic acid* (8.90 g, 43.6 mmol) in water (270 ml) was added NaHCO<sub>3</sub> (7.5 g) and subsequently a solution of benzyl chloroformate (6.85 mL, 48.0 mmol, 1.1 eq.) in Et<sub>2</sub>O (30 mL) was added dropwise with vigorous stirring. Another quantity of NaHCO<sub>3</sub> (7.5 g) was added, and the mixture was stirred vigorously for 3 h. The mixture was washed with Et<sub>2</sub>O (2 x 200 mL), cooled to 0°C, acidified with solid citric acid to pH 3 and extracted with EtOAc (2 x 200 mL). The combined organic phases were washed with 10%
- 20 aq. citric acid (100 mL), dried (Na<sub>2</sub>SO<sub>4</sub>) and concentrated *in vacuo* to a foam. Yield 12.55 g

(85%).  $^1\text{H}$  NMR (acetone- $d_6$ )  $\delta$  11.22 (bs, 1H), 7.40-7.20 (m, 5H), 6.58 (bs, 1NH), 6.23 (d, 1NH,  $J = 7.7$  Hz), 5.07 (s, 2H), 4.35-4.25 (m, 1H), 3.65-3.60 (m, 1H), 3.55-3.50 (m, 1H), 1.39 (s, 9H).  $^{13}\text{C}$ -NMR (acetone- $d_6$ )  $\delta$  171.76, 157.22, 155.89, 137.70, 128.69, 128.11, 78.91, 66.22, 54.56, 42.57, 28.03. MS: expected: 338.4; observed: 338.2.

5

*(2-tert-Butoxycarbonylamino-2-methylcarbamoyl-ethyl)-carbamic acid benzyl ester*

C) To a solution of  $N^{\alpha}$ -Boc- $N^{\beta}$ -Z-Diaminopropionic acid (11.5 g; 34 mmol), HOBt (7.8 g, 51 mmol, 1.5 eq.) and methylamine (8M in ethanol, 6.4 mL, 1.5 eq.) in DCM:DMF (10:1; 110 mL) at 0°C was added DCC (8.4 g, 40.8 mmol, 1.2 eq.) and the mixture was stirred for 2 h while warming to 20°C. 10% aq.  $\text{NaHCO}_3$  (30 mL) was added and the mixture was stirred for further 30 min. The mixture was filtered and the phases separated. The organic phase was extracted with aq.  $\text{KHSO}_4$  (0.5M, 30 mL) and brine (30 mL), dried ( $\text{MgSO}_4$ ) and concentrated *in vacuo* to a white solid. Yield 7.5 g (63%). MS:  $[\text{M}+\text{H}]$  expected: 352.4; observed: 351.9.

*(2-Amino-1-methylcarbamoyl-ethyl)-carbamic acid tert-butyl ester*

D) To a solution of (2-tert-Butoxycarbonylamino-2-methylcarbamoyl-ethyl)-carbamic acid benzyl ester (7.0 g, 19.9 mmol) in methanol (100 mL) was added Pd/C (310 mg). The mixture was hydrogenated for 1 h at 1 atm, filtered (celite) and concentrated *in vacuo* to a foam. Yield 4.3 g (100%). MS:  $[\text{M}+\text{H}]$  expected: 218.30; observed: 218.07.

20

*(2-tert-Butoxycarbonylamino-2-methylcarbamoyl-ethylamino)-acetic acid ethyl ester*

E) To a stirred mixture of (2-Amino-1-methylcarbamoyl-ethyl)-carbamic acid tert-butyl ester (4.2 g, 19.4 mmol) and triethylamine (3.5 mL, 25.2 mmol, 1.3 eq.) in THF (80 mL) was added a solution of ethyl bromoacetate (2.37 mL, 21.3 mmol, 1.1 eq.) in THF (20 mL) dropwise over 15 min. The mixture was stirred for 16 h, filtered and concentrated *in vacuo* to an oil, which was purified by flash chromatography (silica, 7.5% methanol in dichloromethane). Yield: 3.5 g (59%). MS:  $[\text{M}+\text{H}]$  expected: 304.36; observed: 304.04.

*[[2-(6-Benzyloxycarbonylamino-purin-9-yl)-acetyl]-(2-tert-butoxycarbonylamino-2-methylcarbamoyl-ethyl)-amino]-acetic acid ethyl ester*

F) To a stirred suspension of (2-tert-Butoxycarbonylamino-2-methylcarbamoyl-ethylamino)-acetic acid ethyl ester (3.27 g, 10 mmol) and DHBt-OH (1.2 eq.) in DMF (50 mL) was added a solution of (6-benzyloxy-carbonylamino-purin-9-yl)-acetic acid (3.0 g, 10 mmol, 1.0 eq.) in DMF (25 mL). The mixture was cooled to 0°C and DCC (2.26 g, 11 mmol, 1.1 eq.) was added. The mixture was stirred for 1 h at 0°C and then at 20°C for 16 h. DCU was removed

35


by filtration and washed with DMF (20 mL). The filtrate was concentrated in vacuo to 25% volume and DCM (100 mL) was added. The solution was washed with 0.5M KHSO<sub>4</sub> (2 x 50 mL) and 5% aq. NaHCO<sub>3</sub> (2 x 50 mL) and activated carbon (1 g) was added. The mixture was stirred for 1 h, filtered and concentrated in vacuo to an oil, which was purified by flash chromatography (silica, 5% methanol in dichloromethane). Yield: 2.7 g (44%) MS: [M+H]<sup>+</sup> expected: 613.27; observed: 613.10.

*[[2-(6-Benzyloxycarbonylamino-purin-9-yl)-acetyl]-(2-tert-butoxycarbonylamino-2-methylcarbamoyl-ethyl)-amino]-acetic acid*

- G) To a mixture of [[2-(6-Benzyloxycarbonylamino-purin-9-yl)-acetyl]-(2-tert-butoxycarbonylamino-2-methylcarbamoyl-ethyl)-amino]-acetic acid ethyl ester (2.6 g, 4.25 mmol) in THF (50 mL) was added a mixture of LiOH (1.25 eq.) in water (7 mL). The mixture was stirred for 2 h and then concentrated in vacuo to 15% vol. pH was adjusted to ~2.5 with 1M aq. HCl and ethyl acetate (20 mL) was added. The phases were separated and the organic phase was concentrated in vacuo to a foam. Yield: 2.47 g (99%). MS: [M+H]<sup>+</sup> expected: 585.23; observed: 585.3.

### EXAMPLE 53


Compounds of the invention were prepared according to the following reaction schemes. First, a carbohydrate-linker building block is synthesized according to Scheme 1:



**Scheme 1**

Then, a peptide nucleic acid molecule of the invention is synthesized according to Scheme 2:

5


**Scheme 2**

10 wherein B is a naturally-occurring nucleobase preferably A, T, G, or C and n is an integer of from 3 to 49 and wherein the nucleobases are selected in order to bind to the target DNA or target RNA.




**EXAMPLE 54**

Compounds of the invention were prepared as described in Example 53 based upon the following carbohydrate-linker building block:




resulting in a compound of the following formula:




wherein B is a naturally-occurring nucleobase preferably A, T, G, or C and n is an integer of from 3 to 49 and wherein the nucleobases are selected in order to bind to the target DNA or target RNA.

**EXAMPLE 55**

Compounds of the invention were prepared as described in Example 53 based upon the following carbohydrate-linker building block:



resulting in a compound of the following formula:



wherein B is a naturally-occurring nucleobase preferably A, T, G, or C and n is an integer of from 3 to 49 and wherein the nucleobases are selected in order to bind to the target DNA or target RNA.

**EXAMPLE 56 Preparation of GalNAc(OBz)<sub>3</sub>-O-(CH<sub>2</sub>)<sub>4</sub>-COOH (G)****GalNAc(OAc)<sub>4</sub> (A)**

5 D-Galactosamine, HCl (0,185mol; 40g), DMAP (0,018mol; 2,26g) and acetic anhydride (2,4mol; 240mL) were mixed in pyridine (400 mL) at RT. After 16h the reaction mixture was evaporated to an oil, and ethyl acetate (200 mL) was added. The mixture was stirred and after 2 h filtered through a glass filter, and the white product washed with one portion of ethyl acetate. The filtrate evaporated again, and by adding ethyl acetate more product was isolated. The white crystals were dried in vacuum oven at 25°C, 10mbar. Yield 59g (83%)

**Intermediate (B)**

A (0,051mol; 20g) was dissolved in dry 1,2-dichloroethane (250mL) in a dry flask. Trimethylsilyl trifluoromethanesulfonate (0,056mol; 10mL) was added and reacted by heating to 50°C for 2 h. The reaction was performed in N<sub>2</sub>-atmosphere. Triethylamine was added to the reaction until pH 8-9 and the resulting mixture was filtered through a short silica-gel column (2 cm), evaporated and further purified on a silica-gel column using ethyl acetate + 0,1% triethylamine as eluent. Yield: 16,6g (99%)

**GalNAc(OAc)<sub>3</sub>-O-(CH<sub>2</sub>)<sub>5</sub>-OTr (C)**

B (0,050mol; 16,6g) and 5-Trityloxy-pentan-1-ol (0,060mol, 20,8g) were dissolved in dichloroethane (200mL) under N<sub>2</sub>. Molecular sieve 4Å (10g) was added to the reaction. After 30 min trimethylsilyl trifluoromethanesulfonate (0,005mol; 1mL) was added to the yellow mixture. After stirring at room temperature for 20 h, the mixture was added triethylamine (20mL) and filtered through a short silica-gel column (2 cm), The filter was washed 5 times with a mixture of ethyl acetate + 0,1% triethylamine (5 \* 100mL) and the combined organic phases were evaporated. The resulting mixture was purified on a silica-gel column using ethyl acetate/ toluene (6:4) as eluent. Fractions identified as C were collected and evaporated. Yield: 17,4g (50%)

**GalNAc(OBz)<sub>3</sub>-O-(CH<sub>2</sub>)<sub>5</sub>-OTr (D)**

C (0,025mol; 17,2g) was suspended in methanol (220 mL) under N<sub>2</sub> and potassium tert-butoxide (0,002mol; 0,2g) was added. After 1 h the reaction was evaporated to dryness and subsequently treated three times with pyridine (3 \* 30mL) followed by evaporation. The resulting oil was dissolved in pyridine (200mL) under N<sub>2</sub> and benzoic anhydride (0,075mol;

20g) and a catalytic amount of 4-dimethylaminopyridine (0,5g) was added. After stirring at room temperature for 16 h, the mixture was evaporated to dryness, dissolved in ethyl acetate (200mL) and extracted with half-saturated  $\text{NaHCO}_3$  (2 \* 100mL) and water (2 \* 200 mL). DMEDA (20mL) was added to the organic layer, and this was extracted with half-saturated  $\text{KHSO}_4$  (6 \* 100mL), brine (200mL) and dried over  $\text{MgSO}_4$ . Filtering and evaporation gave the product **D**. Yield: 16,4g (75%)

#### **$\text{GalNAc}(\text{OBz})_3\text{-O}-(\text{CH}_2)_5\text{-OH}$ (**E**)**

**D** (0,018mol; 16g) was dissolved in a mixture of methanol/ formic acid (100mL/ 120mL). The mixture was stirred at room temperature for 1 h and slowly added over dry  $\text{K}_2\text{CO}_3$  (approx. 150g). The mixture was stirred in 3 h, and extracted between ethyl acetate (300mL) and water (200mL). The organic layer was drying over  $\text{MgSO}_4$ . Filtering and evaporation gave a crude product, which was further purified on a silica-gel column using toluene/ ethyl acetate (6:4) as eluent. Yield: 8,4g (75%)

#### **$\text{GalNAc}(\text{OBz})_3\text{-O}-(\text{CH}_2)_4\text{-CHO}$ (**F**)**

Oxalylchloride (0,028mol; 2,47mL) was dissolved in DCM (100mL) and cooled on acetone/ dryice bath to  $-60^\circ\text{C}$ . DMSO (30mL) in DCM (2M) was added and reacted in 5 min. A solution of **E** (0,014mol; 8,4g) in DCM (100mL) was added dropwise (10 min). After stirring cold for 40 min triethyl amine (17,3mL) was added and coolbath removed. To the resulting mixture was added ethyl acetate (300mL) and extracted with water (200mL), 10%  $\text{NaHCO}_3$  in water (2 \* 150mL) and finally water (200mL). The organic phase was evaporated to dryness, and subsequently treated three times with toluene followed by evaporation. Yield: 6,8g (99%)

#### **$\text{GalNAc}(\text{OBz})_3\text{-O}-(\text{CH}_2)_4\text{-COOH}$ (**G**)**

**F** (0,011mol; 6,8g) was dissolved in a mixture of  $\text{NaH}_2\text{PO}_4$  (0,073mol; 8,72g),  $\text{NaClO}_2$  (90%)(0,078mol; 8,83g), 2-Methyl-2-butene (0,715mol; 75mL) in acetonitrile/ water (1:1; 400mL). After stirring at room temperature for 1 h, the mixture was added ethyl acetate (50mL) and water (30mL), followed by separation of the organic layer. The aq. phase was extracted with ethyl acetate (3 \* 50mL). The combined organic phases were dried over  $\text{MgSO}_4$ , and finally filtering and evaporation gave the product **G**. LC-MS (electrospray): found  $m/z$  633,2; theoretical  $m/z$  633,2. Purity (RP-HPLC, 251 nm): 97%. Yield: 6,92g (99%).

**EXAMPLE 57 Preparation of GalNAc(OBz)<sub>3</sub>-O-(CH<sub>2</sub>)<sub>4</sub>-CONH-Lys(GalNAc(OBz)<sub>3</sub>-O-(CH<sub>2</sub>)<sub>4</sub>-CONH-)-Gly-OH (K).**

**Boc-Lys(Boc)-Gly-OBn (H)**

- 5 Boc-Lys(Boc)-OSu (11,57mmol; 5,0g) was dissolved in DCM (50mL) and H-Gly-OBzl, Ts (12,4mmol; 4,17g) was dissolved in DCM, deprotonated with DIEA (12,4mmol; 2,0mL) and added to the reaction. The mixture was stirred for 1 h, extracted with 50% sat. aq. NaHCO<sub>3</sub> (50mL) and 50% sat. aq. KHSO<sub>4</sub> (50mL). The organic phase was dried over MgSO<sub>4</sub> and concentrated *in vacuo* to an oil. Yield: 5,5g (99%).

10

**GalNAc(OBz)<sub>3</sub>-O-(CH<sub>2</sub>)<sub>4</sub>-CONH-Lys(GalNAc(OBz)<sub>3</sub>-O-(CH<sub>2</sub>)<sub>4</sub>-CONH-)-Gly-OBn (J)**

HOAt (1,32mmol; 180mg), EDC (1,32mmol; 254mg), DIEA (1,32mmol; 230μL) and G (1,2mmol; 543mg) were dissolved in dry DCM (4mL). Reaction time 3 h.

- 15 H (0,551mmol; 272mg) was stirred in DCM/ trifluoroacetic acid (1:1; 6mL) for 1 h and then concentrated *in vacuo* and coevapored with DCM (5mL), toluene (5mL) and DCM (2 \* 5mL). The resulting oil was dissolved in DCM/ DMF (5:1; 6mL). pH adjust to 8 with NMM and added to the mixture of the activated acid. Reaction time 20 h. Ethyl acetate (8mL) and water (5mL) was added to the reaction, followed by separation of the organic layer. The organic
- 20 layer was extracted with 50% sat. aq. KHSO<sub>4</sub> (2 \* 5mL), 50% sat. aq. NaHCO<sub>3</sub> (2 \* 5mL), water (2 \* 5mL) and brine (5mL). The organic phase was dried over MgSO<sub>4</sub>, filtered and evaporated to give the crude product, which was purified on a silica-gel column using DCM/ methanol (15:1) as eluent. Yield: 420mg (50%).

- 25 **GalNAc(OBz)<sub>3</sub>-O-(CH<sub>2</sub>)<sub>4</sub>-CONH-Lys(GalNAc(OBz)<sub>3</sub>-O-(CH<sub>2</sub>)<sub>4</sub>-CONH-)-Gly-OH (K)**

J (0,275mmol; 420mg) was dissolved in ethanol/ THF (1:1; 75mL) under N<sub>2</sub>, 10% Pd/C (50mg) was added and the mixture was hydrogenated with H<sub>2</sub> gas. After 2 h the reaction mixture was filtered through Hyflo®, which was rinsed afterwards with 1 portion of ethyl acetate (20mL). The combined organic phases were evaporated to dryness, and subsequently

30 treated three times with toluene followed by evaporation. LC-MS (electrospray): found m/z 1434,3; theoretical m/z 1433,6. Purity (RP-HPLC, 210 nm): 97%. Yield: 395mg (99%).

**EXAMPLE 58 Preparation of GalNAc(OBz)<sub>3</sub>-O-(CH<sub>2</sub>)<sub>4</sub>-CONH-Lys(GalNAc(OBz)<sub>3</sub>-O-(CH<sub>2</sub>)<sub>4</sub>-CONH)-Lys(GalNAc(OBz)<sub>3</sub>-O-(CH<sub>2</sub>)<sub>4</sub>-CONH)-Gly-OH (S)**

35

**Boc-Lys(Z)-OMe (L)**

Boc-Lys(Z)-OH (0,026mol; 10,0g) and NaHCO<sub>3</sub> (0,079mol; 6,6g) were dissolved in DMF (75mL) in a dark flask (light sensitive!). Methyl iodide (0,029mol; 1,8mL) was added to the reaction. After stirring at room temperature for 16 h, ethyl acetate (400mL) was added and the mixture was extracted with water (2 \* 100mL), brine (100mL), dried over MgSO<sub>4</sub> and filtered through a short silica-gel column. The filtrate was concentrated *in vacuo* to a yellow oil. Yield: 10,2g (99%).

**Boc-Lys-OMe (M)**

L (0,028mol; 11,6g) was dissolved in a mixture of ethanol/ THF (1:1; 100mL) under N<sub>2</sub>. 10 % Pd/C (0,6g) was added and the mixture was hydrogenated with H<sub>2</sub> gas. After 1 h the reaction mixture was filtered through Hyflo®, which was rinsed afterwards with 2 portions of ethyl acetate (2 \* 25mL). The combined organic phases were evaporated to dryness, and subsequently treated three times with toluene followed by evaporation. Yield: 7,9g colourless oil (99%).

**Boc-Lys(Boc-Lys(Boc))-OMe (N)**

M (0,007mol; 1,85g) was dissolved in dry DCM (20mL) under N<sub>2</sub>. After cooling to 0°C on ice bath Boc-Lys(Boc)-OSu (0,007mol; 3,2g) was added to the reaction and the ice bath was removed. The mixture was stirred for 5 h. DMEDA (2mL) was added. After stirring for 15 min the reaction mixture was extracted with water (20mL), 20% KHSO<sub>4</sub> (20mL), water (20mL) and brine (20mL). The organic layer was dried over MgSO<sub>4</sub>, filtered and evaporated. The resulting mixture was purified on a silica-gel column using DCM/ methanol (19:1) as eluent. Yield: 2,7g white crystals (65%).

**Boc-Lys(Boc-Lys(Boc))-OH (P)**

N (0,004mol; 2,7g) was dissolved in THF (200mL) and 1N LiOH (15mL) was added. After stirring for 3 h water (100mL) was added, and pH adjusted with 2N HCl to pH 3. The mixture was extracted with ethyl acetate (3 \* 100mL) and the combined organic phases were extracted with brine (150mL), dried over MgSO<sub>4</sub>, filtered and evaporated to dryness. Yield: 2,65g colourless oil (99%).

**Boc-Lys(Boc-Lys(Boc))-Gly-OBn (Q)**

**P** (0,0045mol; 2,6g) was dissolved in DCM (50mL) and cooled to -10°C. H-Gly-OBzl, **Ts** (0,0054mol; 1,82g) was deprotonated with DIEA and added to the reaction. HOBt (0,0054; 0,73g), EDC (0,0054mol; 1,03g) were added and the pH was adjusted with DIEA to pH 7-8. After 2 h reaction pH was adjusted again. After 20 h reaction 10% aq. NaHCO<sub>3</sub> (30mL) was added. After stirring for 30 min the reaction was extracted with water (2 \* 50mL), 50% sat. aq. KHSO<sub>4</sub> (2 \* 50mL), water (2 \* 50 mL) and brine (50mL). The organic phase was evaporated, and the resulting oil was purified on a silica-gel column using DCM/ methanol (19:1) as eluent. Yield: 2,2g colourless oil (68%).

- 10 **GalNAc(Obz)<sub>3</sub>-O-(CH<sub>2</sub>)<sub>4</sub>-CONH-Lys(GalNAc(Obz)<sub>3</sub>-O-(CH<sub>2</sub>)<sub>4</sub>-CONH)-Lys(GalNAc(Obz)<sub>3</sub>-O-(CH<sub>2</sub>)<sub>4</sub>-CONH)-Gly-OBn (**R**)**  
 HOAt (1,0mmol; 0,14g), EDC (1,1mmol; 0,21g), DIEA (1,2mmol; 210μL) and **G** (1,0mmol; 0,64g) were dissolved in dry DCM (10mL). Reaction time 3 h. In another flask **Q** (0,305mmol; 0,22g) was dissolved in 5mL DCM and trifluoroacetic acid (5mL) was added. After 2 h reaction the mixture was evaporated and pH adjust to 8-9 with DIEA. The deprotected compound **Q** was dissolved in DCM (5mL) and DMF (1mL) and added to the mixture of the activated acid. Reaction time 20 h. Ethyl acetate (20mL) and water (25mL) was added to the reaction, followed by separation of the organic layer. The organic layer was extracted with 50% sat. aq. KHSO<sub>4</sub> (2 \* 20mL), 50% sat. aq. NaHCO<sub>3</sub> (2 \* 20 mL), water (2 \* 20mL) and brine (20mL). The organic phase was dried over MgSO<sub>4</sub>, filtered and evaporated to give the crude product, which was purified on a silica-gel column using DCM/ methanol (15:1) as eluent. Yield: 76mg colourless oil (50%).

- 25 **GalNAc(Obz)<sub>3</sub>-O-(CH<sub>2</sub>)<sub>4</sub>-CONH-Lys(GalNAc(Obz)<sub>3</sub>-O-(CH<sub>2</sub>)<sub>4</sub>-CONH)-Lys(GalNAc(Obz)<sub>3</sub>-O-(CH<sub>2</sub>)<sub>4</sub>-CONH)-Gly-OH (**S**)**  
**R** (0,003mmol; 76mg) was dissolved in ethanol/ THF (1:1; 10mL) under N<sub>2</sub>, 10% Pd/C (25mg) was added and the mixture was hydrogenated with H<sub>2</sub> gas. After 2 h the reaction mixture was filtered through Hyflo®, which was rinsed afterwards with 1 portion of ethyl acetate (4mL). The combined organic phases were evaporated to dryness, and subsequently treated three times with toluene followed by evaporation. LC-MS (electrospray): found m/z 2176,9; theoretical m/z 2177,3. Purity (RP-HPLC, 254 nm): 97%. Yield: 60 mg colourless oil (82%).

- 35 **EXAMPLE 59 Preparation of GalNAc(Obz)<sub>3</sub>-O-(CH<sub>2</sub>)<sub>4</sub>-CONH -Lys(GalNAc(Obz)<sub>3</sub>-O-(CH<sub>2</sub>)<sub>4</sub>-CONH -Lys(GalNAc(Obz)<sub>3</sub>-O-(CH<sub>2</sub>)<sub>4</sub>-CONH -Lys(GalNAc(Obz)<sub>3</sub>-O-(CH<sub>2</sub>)<sub>4</sub>-CONH)))-**

**Gly-OH (Z)****Boc-Lys(Boc-Lys(Z))-OMe (T)**

**T** (18,05mmol; 4,7g) was dissolved in dry DCM/ dry DMF (10:1; 100mL) under N<sub>2</sub>. After cooling to 0°C on ice bath Boc-Lys(Z)-OH (18,05mmol; 6,9g), HOBt (18,05mmol; 2,4g), EDC (19,85mmol; 3,8g) and DIPEA (27mmol; 4,7mL) was added to the reaction and the ice bath was removed. The mixture was stirred for 20 h. The reaction mixture was extracted with water (100mL), 20% sat. aq. KHSO<sub>4</sub> (100mL), water (100mL) and brine (100mL). The organic layer was dried over MgSO<sub>4</sub>, filtered and evaporated. The resulting oil was precipitated with DCM/ Petroleum ether (40-60°C). Yield: 6,7g white crystals (60%).

**Boc-Lys(Boc-Lys(H))-OMe (U)**

**U** (10,44mmol; 6,5g) was dissolved in a mixture of ethanol/ THF (1:1; 100mL) under N<sub>2</sub>. 10 % Pd/C (0,5g) was added and the mixture was hydrogenated with H<sub>2</sub> gas. After 2 h the reaction mixture was filtered through Hyflo®, which was rinsed afterwards with 2 portions of ethyl acetate (2 \* 50mL). The combined organic phases were evaporated to dryness, and subsequently treated three times with toluene followed by evaporation. Yield: 5,1 g colourless oil (99%).

**Boc-Lys(Boc-Lys(Boc-Lys(Boc)))-OMe (V)**

**V** (10,64mmol; 5,1g) was dissolved in dry DCM (20mL) under N<sub>2</sub>. After cooling to 0°C on ice bath Boc-Lys(Boc)-OSu (10,64; 4,7g) was added to the reaction and the ice bath was removed. The mixture was stirred for 20 h. The reaction mixture was extracted with water (20mL), 20% KHSO<sub>4</sub> (20mL), water (20mL) and brine (20mL). The organic layer was dried over MgSO<sub>4</sub>, filtered and evaporated. The resulting mixture was purified on a silica-gel column using DCM/ methanol (15:1) as eluent. The resulting oil was dissolved in hot ethyl acetate and precipitated with hexane. Yield: 4,7g white crystals (55%).

**Boc-Lys(Boc-Lys(Boc-Lys(Boc)))-OH (W)**

**W** (4,89mmol; 4,0g) was dissolved in THF (60mL) and 1N LiOH (7,3mL) was added. After stirring for 6 h pH adjusted with 2N HCl to pH 4. The mixture was evaporated to an oil, dissolved in DCM (50mL) and extracted with water (50mL), brine (150mL), dried over MgSO<sub>4</sub>, filtered and evaporated to an oil. Yield: 3,95g colourless oil (99%).



**Boc-Lys(Boc-Lys(Boc-Lys(Boc)))-Gly-OBn (X)**

**W** (4,89mmol; 4g) was dissolved in DCM/ DMF (5:1; 50mL) and cooled to -10°C. H-Gly-OBzl, p-TsOH (4,89mmol; 1,7g) was dissolved in DCM, deprotonated with DIEA (12,25mmol; 2,1mL) and added to the reaction. HOBt (4,89mmol; 0,66g), EDC (5,38mmol; 1,03g) were added and the pH was adjusted with DIEA to pH 7-8. After 2 h reaction pH was adjusted again. After 20 h reaction 10% aq. NaHCO<sub>3</sub> (30mL) was added. After stirring for 30 min the reaction was extracted with water (2 \* 50mL), 50% sat. aq. KHSO<sub>4</sub> (2 \* 50mL), water (2 \* 50 mL) and brine (50mL). The organic phase was evaporated, and the resulting oil was dissolved in refluxing ethyl acetate. The mixture was placed on an ice bath and the compound precipitate after 20min. Yield: 4,1g white crystals (87%).

**GalNAc(Obz)<sub>3</sub>-O-(CH<sub>2</sub>)<sub>4</sub>-CONH -Lys(GalNAc(Obz)<sub>3</sub>-O-(CH<sub>2</sub>)<sub>4</sub>-CONH -Lys(GalNAc(Obz)<sub>3</sub>-O-(CH<sub>2</sub>)<sub>4</sub>-CONH -Lys(GalNAc(Obz)<sub>3</sub>-O-(CH<sub>2</sub>)<sub>4</sub>-CONH))) -Gly-OBn (Y)**

HOAt (6,94mmol; 0,88g), EDC (7,57mmol; 1,31g), DIEA (9,47mmol; 1,31mL) and **G** (6,313mmol; 4,0g) were dissolved in dry DCM (50mL). Reaction time 3 h. In another flask **X** (1,05mmol; 1,0g) was dissolved in DCM (10mL) and trifluoroacetic acid (10mL) was added. After 2 h reaction the mixture was evaporated and pH adjust to 8-9 with DIEA. The deprotected compound **X** was dissolved in DCM (50mL) and DMF (20mL) and added to the mixture of the activated acid. Reaction time 20 h. Ethyl acetate (200mL) and water (100mL) was added to the reaction, followed by separation of the organic layer. The organic layer was extracted with 50% sat. aq. KHSO<sub>4</sub> (50mL), 50% sat. aq. NaHCO<sub>3</sub> (50mL), water (50mL) and brine (50mL). The organic phase was dried over MgSO<sub>4</sub>, filtered and evaporated to give the crude product, which was purified on a silica-gel column using DCM/ methanol (15:1) as eluent. Yield: 1,5g (40%).

**GalNAc(Obz)<sub>3</sub>-O-(CH<sub>2</sub>)<sub>4</sub>-CONH -Lys(GalNAc(Obz)<sub>3</sub>-O-(CH<sub>2</sub>)<sub>4</sub>-CONH -Lys(GalNAc(Obz)<sub>3</sub>-O-(CH<sub>2</sub>)<sub>4</sub>-CONH -Lys(GalNAc(Obz)<sub>3</sub>-O-(CH<sub>2</sub>)<sub>4</sub>-CONH))) -Gly-OH (Z)**

**Y** (0,498mmol; 1,5g) was dissolved in ethanol/ THF (3:1; 100mL) under N<sub>2</sub>, 10% Pd/C (150mg) was added and the mixture was hydrogenated with H<sub>2</sub> gas (balloon). After 20 h the reaction mixture was filtered through Hyflo®, which was rinsed afterwards with 1 portion of ethyl acetate (100mL). The combined organic phases were evaporated to dryness, and subsequently treated three times with toluene followed by evaporation. LC-MS (electrospray): found m/z 2921,1; theoretical m/z 2920,2. Purity (RP-HPLC, 254 nm): 91%. Yield: 1,2g (82%).

**EXAMPLE 60 Preparation of (GalNAc(OH)<sub>3</sub>)<sub>2</sub>-Lys-Gly-CATCACTGGCAGACCCTG-NH<sub>2</sub>**

5 In a 10-mL round-bottomed flask an aeg-PNA of the sequence H-CATCACTGGCAGACCCTG-NH<sub>2</sub> (85,42 mg; 0,0176 mmoles) and K (50,60 mg; 0,0353 mmoles, 2,00 eq.) were dissolved in DMSO (7,7 mL). A 0,125M solution of HOBt in DMSO (296,3 µL; 0,0370 mmoles, 2,10 eq.) was added and pH (5-6) adjusted to 7-8 by addition of 10% NMM in DMSO (500 µL). A 0,125M solution of DIC in DMSO (564,4 µL; 0,0706 mmoles, 4,00 eq.) was finally added and the reaction mixture was stirred at room temperature for 45 h.

Deprotection of the benzoyl groups was performed by addition of 1% NaOH in MeOH (1,47 mL) to the reaction mixture until the pH was raised to 10-12. The debenzoylation reaction was completed after 2h 30min. and the crude construct was precipitated by pouring the mixture over cold diethyl ether (90 mL). The suspension thus obtained was centrifuged, the ether discarded and the pellet washed thoroughly two times with ether (2 x 90 mL). The resulting pellet was dissolved in water (11mL) and the pH adjusted to 1-2 with TFA to facilitate dissolution of the solid.

The crude solution was purified by RP-HPLC and after freeze-drying 21 mg of the pure compound was obtained as a white powder. LC-MS (electrospray): found *m/z* 5631,8; theoretical *m/z* 5632,5. Purity (RP-HPLC, 210 nm): 97%.

Total yield: 21,1%

**EXAMPLE 61 Preparation of (GalNAc(OH)<sub>3</sub>)<sub>3</sub>Lys<sub>2</sub>Gly-GTGGATGATACCTGGATC-NH<sub>2</sub>**

10 In a 25-mL round-bottomed flask an aeg-PNA of the sequence GTGGATGATACCTGGATC-NH<sub>2</sub> (200,62 mg; 0,04052 mmoles) was dissolved in DMSO (4 mL). A solution of S (140,92 mg; 0,06469 mmoles, 1,60 eq) in DMSO (1 mL) was added followed by a 0,125M solution of HOBt in DMSO (543 µL; 0,06787 mmoles, 1,68 eq). The pH (≈3) was raised to 7-8 by addition of 10% NMM in DMSO (1,1 mL). Then, a 0,125M solution of DIC in DMSO (2,07 mL; 0,2588 mmoles, 6.40 eq) was added together with DMSO (1,29 mL) to reach a final concentration of 20 mg PNA/mL. The reaction mixture was stirred at room temperature for 21 h 30 min.

35 To perform debenzoylation, 1%NaOH in MeOH was added over the mixture until de pH was

raised to 10-12 (3,2 mL). After a few seconds the solution turned opalescent and afterwards a fine white powder precipitated. Deprotection of the benzoyl groups was completed after 2 h. The crude was precipitated in cold diethyl ether (160 mL), pelleted by centrifugation and further washed with ether (200 mL). The crude compound was isolated by centrifugation,  
5 dissolved in water (6mL) and pH adjusted to 2-3.

The crude solution was purified by RP-HPLC and after freeze-drying, 140,7 mg of pure compound was obtained as a white powder. LC-MS (electrospray): found  $m/z$  6173,8; theoretical  $m/z$  6174,1. Purity (RP-HPLC, 210 nm): 96%.

Total yield: 56,2%

10

#### **EXAMPLE 62 Preparation of (GalNAc(OH)<sub>3</sub>)<sub>4</sub>Lys<sub>3</sub>Gly-GTGGATGATACCTGGATC-NH<sub>2</sub>**

In a 25-mL round-bottomed flask, an aeg-PNA of the sequence H-

15 GTGGATGATACCTGGATC-NH<sub>2</sub> (350,42 mg; 0,0708 mmoles) was dissolved in DMSO (5 mL). A solution of Z (333,50 mg; 0,1141 mmoles, 1,61 eq) in DMSO (2 mL) was added followed by a 0,125M solution of HOBt in DMSO (951  $\mu$ L; 0,1188 mmoles, 1,68 eq). By addition of 10% NMM in DMSO (1,8 mL), the pH ( $\approx$ 3) was raised to 7-8 and then a 0,125M solution of DIC in DMSO (3,62 mL; 0,4525 mmoles, 6.39 eq) was added together with DMSO (4,13 mL)  
20 to reach a final concentration of 20 mg PNA/mL. The reaction mixture was stirred at room temperature for 21h 30'.

Debenzoylation was performed by addition of 1% NaOH in MeOH (7,5mL) until the pH was raised to 10-12. A fine white suspension was obtained, and after 2h 40min. the reaction was completed. The mixture was poured over cold diethyl ether (160 mL), the compound separated by centrifugation, washed thoroughly with ether (200 mL) and the suspension further  
25 centrifuged. The crude pellet was dissolved in water (12 mL) and the pH adjusted to 2-3.

After purification (RP-HPLC) and freeze-drying, 200,08 mg of compound was obtained as a white powder. LC-MS (electrospray): found  $m/z$  6604,5; theoretical  $m/z$  6605,6. Purity (RP-HPLC, 210 nm): 96%.

30 Total yield: 42,8%.

#### **EXAMPLE 63 Preparation of Folate-CCTCTTACCTCAGTTACA-NH<sub>2</sub>**

35 An aeg-PNA of the sequence H-CCTCTTACCTCAGTTACA-NH<sub>2</sub> (10,00 mg; 0,0021mmoles)

was dissolved in DMSO (1 mL). A 0,13M solution of folic acid in DMSO (161,4  $\mu$ L; 0,0210 mmoles) was added together with a 0,125M solution of HOBt in DMSO (167,8  $\mu$ L; 0,0210 mmoles). The pH of the mixture (5-6) was adjusted to 7-8 by addition of 10% NMM in DMSO (60  $\mu$ L), and then a 0,125M solution of DIC in DMSO (167,8  $\mu$ L; 0,0210 mmoles) was added.

- 5 The mixture was stirred at room temperature for 21h 30 min. The conjugate and excess of folic acid were precipitated by pouring the reaction mixture over cold diethyl ether (10 mL) and pelleted by centrifugation. The pellet was washed twice with ether (2 x10 mL) and then dissolved in a mixture of DMSO (0,5 mL) and 0,1% TFA in water (3,5mL).  
After purification (RP-HPLC) and freeze-drying, 5,00 mg of compound was obtained as a yellow powder. LC-MS (electrospray): found m/z 5188,1; theoretical m/z 5189,1. Purity (RP-HPLC, 210 nm): 94%  
Total yield: 45,9%

#### 15 **EXAMPLE 64 Preparation of PNA and PNA Conjugates**

- The PNA was synthesized on MBHA-resin in a sequential manner. The oligomers were synthesized from Boc-protected PNA-monomers using a standard SPPS protocol. The products were cleaved from the resin using trifluoromethane sulphonic acid. Crude deprotected PNAs  
20 were purified by preparative HPLC and subsequently lyophilized.

Two 18mer PNA oligomers, PNA-I and PNA-II, were synthesized, cleaved, purified and lyophilized. The compounds were isolated in 1 g quantities.

#### 25 **GalNAc constructs**

GalNAc(OBz)<sub>3</sub>-O-pentanoic acid (A, Figure 1) was prepared in five steps according to known procedures. The synthesis, which starts from inexpensive D-galactosamine, is high yielding and scalable.




Figure 1

A was used in the preparation of di-, tri- and tetra-antennary ligands (Figure 2-4). Solid-phase synthesis of similar di- and tri-antennary lysine-based cluster galactosides has previously been reported. However, attempts to apply the described procedures resulted in very poor yields and, consequently, the synthetic protocol was redesigned for solution phase synthesis. In contrast to the original solid phase procedure the developed procedure is robust and scalable.

Achiral glycine was incorporated as a C-terminal linker in the cluster galactosides to avoid racemization in the subsequent condensation with the PNA oligomer.

The di-antennary ligand (C, Figure 2) was synthesized by initial coupling of pre-activated Boc-Lys(Boc)-OSu to H-Gly-OBn giving scaffold Boc-Lys(Boc)-Gly-OBn (B). Cleavage of the Boc groups and coupling with IBC activated GalNAc(OBz)<sub>3</sub>-O-pentanoic acid (A) gave the di-substituted construct, which was hydrogenated to the desired product (C).




Figure 2

In the synthesis of the tri- and tetra-antennary GalNAc-constructs three orthogonal protecting groups were necessary and, hence, the synthesis was performed in a slightly different man-

ner. Boc-Lys(Z)-OH was initially methylated and the Z-group was removed by hydrogenation giving Boc-Lys-OMe (D). Coupling with pre-activated Boc-Lys(Boc)-OSu gave Boc-Lys(Boc-Lys(Boc))-OMe (E). Hydrolysis of the methyl ester and subsequent coupling with H-Gly-OBn gave Boc-Lys(Boc-Lys(Boc))-Gly-OBn (F). Removal of the Boc groups and coupling with IBC-activated GalNAc(OBz)<sub>3</sub>-O-pentanoic acid (A) gave the tri-substituted construct, which was hydrogenated yielding the desired product (G).




Figure 3

The tetra-antennary GalNAc-construct (H, Figure 4) was synthesized in a manner similar to the tri-antennary GalNAc-construct (G)

70




Figure 4

- 5 The GalNAc monomer (A) was synthesized in 15 g scale, and the three ligands were synthesized in 2-5 g quantities.

#### Ligand-PNA Conjugation

- Previously, transporter-PNA constructs have been synthesized in a sequential manner, e.g. KFF-PNA, or by different cross-linking methods involving thiols, e.g. disulfides or maleimides. These methodologies are not compatible with the GalNAc-PNA chemistry for a number of reasons. The sequential synthesis involves coupling of the GalNAc-construct to PNA on resin. It is not feasible because of the subsequent cleavage conditions, which will also cleave the glycoside bonds. Disulfides might be attractive for screening purposes, but the poor biological and chemical stability of the disulfide bond impedes the use in a ligand-S-S-PNA drug. Incorporation of maleimides, e.g. using SMCC, is unattractive because of the resulting introduction of the bulky linker between the ligand and the PNA. Furthermore, the maleimide coupling results in two diastereomers.

- 20 As a consequence of the limitations in the known cross-linking methods the development of an alternative procedure was initiated. It was attempted to perform a solution phase segment condensation involving an amid bond formation between the C-terminal of the GalNAc-construct and the N-terminal of the PNA.

- 25 There are numerous factors that complicate such a condensation. The poor solubility of PNA in common organic solvents limits the choice of solvent to DMSO. The degree of activation on the carboxylic acid was limited by the demand of high selectivity between the N-terminal

amine and the amines on the nucleobases. Furthermore, a quantitative and clean de-protection of the benzoyl protecting groups on the GalNAc-moieties was mandatory. Finally, the product should be isolated from the DMSO and dissolved in an aqueous buffer for purification.

5

A clean and high yielding one-pot segment condensation/de-protection was developed. The synthesis was performed at pH 7-8 in DMSO using DIC/HOBt activation. The subsequent hydrolysis of the Benzoyl protecting groups was achieved by addition of NaOH in methanol. Finally, the product was precipitated from the reaction mixture in diethyl ether.

10

The method is versatile and does not limit the choice of linker. It allows the incorporation of any ligand with a free carboxylic acid, e.g. protected peptides, bile acids and folic acid. Furthermore the formed amid bond is chemically and enzymatically stable.

15 Four constructs were synthesized in 100 mg scale:

- Construct Ia, based upon PNA-I and a ligand of formula **C**
- Construct Ib, based upon PNA-I and a ligand of formula **G**
- Construct IIa, based upon PNA-II and a ligand of formula **G**
- Construct IIb, based upon PNA-II and a ligand of formula **H**.

20

PNA-I is an aeg-PNA of the sequence H-CATCACTGGCAGACCCTG-NH<sub>2</sub>

PNA-II is an aeg-PNA of the sequence H-GTGGATGATACCTGGATC-NH<sub>2</sub>

25 Synthesized compounds were purified on an automated preparative HPLC system and subsequently lyophilised. Identity and purity were determined using analytical HPLC and mass spectrometry. All compounds had an HPLC purity &gt;95% at 210 nm.

30 **EXAMPLE 65 In vivo testing of PNA and PNA Conjugates**

*Identification of PNAs provoking exon skipping of pre-mRNA expressed in mouse liver*

..

35 Classical antisense molecules recruit RNaseH activity, which leads to degradation of the targeted mRNA, which in turn leads to reduced synthesis of the protein encoded by the targeted



mRNA. This can be visualized by Northern blot, quantitative RT-PCR and Western blot analysis. However there is always a risk that such down regulations are due to unspecific effects on gene expression or translation not related to the desired antisense activity on the intended target. For this reason the lack of effect of scrambled or mismatch oligonucleotides is frequently reported in parallel. However such compounds are poor controls since alterations in the nucleotide sequence is likely to affect the compounds overall pharmaco-kinetic properties.

We therefore sought to obtain compounds with an inbuilt antisense control. We chose to target the splice pattern of pre-mRNA. There are numerous examples in the literature of non-RNaseH-recruiting oligonucleotides capable of changing the splice pattern of pre-mRNA by hybridising to and thereby blocking a splice site. Such effects are easily visualized by RT-PCR analysis. It is very unlikely that splice variations specific for the targeted exon could occur by unspecific effects not related to a direct antisense interaction at the intended target.

#### *Hnf4*

The *Hnf4* gene, encoding the Hepatocyte Nuclear Factor 4 $\alpha$  (HNF4  $\alpha$ ), was chosen as a model target for PNA liver-uptake studies for a number of reasons:

1. It is expressed in hepatocytes carrying the asialoglycoprotein (ASGP) receptors
2. it encodes a transcription factor of which both mRNA and protein are expected to have a short half-life
3. it is constitutively expressed, enabling simple experimental testing

PNAs were designed as 18-mers, spanning the Intron/exon or exon/intron boundary with 9 bases overlapping either side. The PNA sequence was checked for possible problematic characteristics (self-complementarity, high purine content and long N-terminal purine stretches). Possible problems were corrected by choosing adjacent 18-mer sequences (still overlapping the splice site, but non-symmetrical) free of unwanted characteristics. A total of 13 PNAs were designed targeting different splice sites of *Hnf4* pre-mRNA. The PNAs were evaluated *in vitro* using a BNL CL.2 mouse liver cell line. PNA intracellular delivery was achieved by annealing the PNA to a partly complementary DNA-oligo followed by co-transfection using a polyethylenimine transfection reagent.

PNA effect on the splice pattern of *Hnf4* mRNA was evaluated by RT-PCR on RNA extracted

20 hrs. after PNA-treatment of the cells.

Among active PNAs, PNA-II (H-GTGGATGATACCTGGATC-NH<sub>2</sub>) and PNA-I (H-CATCACTGGCAGACCCTG-NH<sub>2</sub>) – were selected for *in vivo* studies.

5

PNA-II (targeting the 3'-end of exon 7) causes skipping exclusively of Hnf4 exon 7, observed as a new shorter RT-PCR product obtained with PCR primers external to exon 7. Minute hardly detectable levels of this splice variant does occur naturally, but the ratio between the normal and the rare splice variant is strongly shifted following PNA-II treatment.

10 PNA-II binds to Hnf4 pre-mRNA with all 18 bases and with 7 bases to correct spliced Hnf4 mRNA.

PNA-I (targeting the 5'-end of exon 9) causes exclusive skipping of Hnf4 exon 9. Again, very low levels of this splice variant occurs naturally, but the ratio between the normal and the rare splice variant, was strongly shifted following PNA-I treatment. PNA-I binds to Hnf4 pre-mRNA with all 18-bases and with coincidently with no less than 17 bases to correct spliced Hnf4 mRNA. As a consequence the PNA is capable of binding to Hnf4 mRNA and inhibits reverse transcriptase. Thus, traditional RT-PCR could not be used for evaluating the effect on splicing. However this problem was solved with a specially designed PCR triple primer mix (see "Analysis of liver samples" below).

Exon 7 and exon 9 skipping both yield *in frame* deletions in the reading frame of hnf4 mRNA. As a result a shorter HNF4 protein was synthesized which could be detected by Western blot analysis. This feature is an additional internal control of antisense activity and made PNA-II and PNA-I particularly interesting candidates for GalNAc modification.

### *Animal studies*

#### *Animals and housing*

30 NMRI female mice (approx. weight 25 to 30 g at the day of the study) were housed in transparent macrolone cages and light cycles of 12 hours light and 12 hours darkness were used. The room temperature was 21°C ± 1°C and the relative humidity 50±10%. The animals were fed ad libitum using a complete rodent diet and had free access to clean drinking water. On the day of study the animals were randomly allocated to groups of two animals each for the different treatments.

35

### *Experimental design*

Animals were dosed subcutaneously (s.c.). For dosing, 400  $\mu$ M solutions (as determined by spectrometry from a theoretically calculated coefficient of extinction) of the respective test compounds in 0.625 mM AcOH, 5 % glucose in water, were used.

At each dosing, the animal received 0.3 ml of the respective PA. Each dose corresponded to approximately 4.8  $\mu$ mol PNA/kg equivalent to approximately 25 mg/kg. For multiple dosings, 6-hour intervals were used between same day dosings and 18-hour intervals before next day dosing.

### *Sampling*

At the indicated number of days after the first dosing, the animals were anaesthetized by carbon di-oxide/oxygen and euthanized by partial decapitation and exsanguination. Immediately hereafter, the abdominal cavity was opened and the liver excised. From the liver, the caudate lobe was isolated (approximate weight 0.1 g) and transferred to 1 ml of RNA preservation solution (RNAlater). Liver tissue for Western blot analysis was immersed in a protease inhibitor solution (ProPrep), homogenized and frozen for later analysis. The remaining liver tissue was frozen and stored at  $-18^{\circ}\text{C}$  for later compound analysis (not presented here).

From selected animals, different other tissues were also excised for RT-PCR analysis for Hnf4: a 3 mm section from the middle of the sagittally cut left kidney, a 3 mm transverse section from the middle of colon descendens, a 3 mm transverse section from the middle of jejunum, and a 3 mm transverse section from the pancreas. These tissues were preserved by immersion in 1 ml of RNA preservation solution (RNAlater).

### *Analysis of liver samples*

#### *RT-PCR*

Approximately 15-20 mg the stabilized liver sample was transferred into 1,0 ml RTL lysis buffer and homogenized. Total RNA was purified according to the RNeasy mini kit protocol and the concentration determined by OD260 measurement.

cDNA was synthesized from 1  $\mu$ g of RNA using random decamer primers in a total vol. of 20  $\mu$ l. Briefly, RNA was preincubated at  $70^{\circ}\text{C}$  for 4 min prior to addition of M-MLV reverse transcriptase, followed by 60 min incubation at  $42^{\circ}\text{C}$  in the presence of 0,5 mM of each dNTP.

The PCR reaction was carried out in 35 cycles using SuperTaq polymerase in presence of

0,4  $\mu$ M primers and 0,2 mM of each dNTP.

For PNA-II a single set of primers could be used as this sequence does not inhibit reverse transcription of Hnf4 mRNA:

caatgaatatgcctgcctcaa (forward)

5 attcagatcccagaccactt (reverse)

For PNA-I, RT-PCR analysis was complicated by the fact that PNA-I is a highly potent inhibitor of reverse transcription of Hnf4 mRNA between exon9 and exon8 due to extensive base complementarity to the exon8/9 junction. This effect is specific for normally spliced Hnf4 mRNA. PNA-I does not affect reverse transcription of exon9 skipped Hnf4 mRNA. It has been found that PNA contaminates the RNA extractions from liver tissue. As a consequence cDNA formation of normally spliced mRNA is selectively inhibited at exon9 and the result of the ensuing PCR reaction (with primers surrounding exon 9) is therefore artificially skewed in favour of the exon9-skipped cDNA.

Exon 9 skip caused by PNA-I was therefore evaluated using a modified RT-PCR strategy based on two independent PCR reactions that specifically amplify either normally spliced or exon9 skipped Hnf4 mRNA. The two reactions were performed simultaneously in the same tube using a triple primer set. The forward primer in exon 10 is common to the two reactions. Two different reverse primers were designed to hybridise specifically with the exon8/10 junction (in exon 9 skipped mRNA) and specifically with exon 9 respectively (in the normally spliced mRNA). The two reverse primers were in addition such designed that they give rise to slightly differently sized PCR products that can be separated and distinguished on an agarose gel. To validate this concept, PCR products specific for normally spliced mRNA and exon9 skipped mRNA were generated and mixed in 100:1, 10:1, 1:1, 1:10 and 1:100 relative concentrations. Subsequent PCR with the triple primer set yielded two products of the expected sizes that accurately reported the relative concentration of template (as long as the templates were sufficiently diluted). The technique was subsequently successfully used to evaluate the efficacy of PNA-I.

The triple primer set used for RT-PCR of mRNA from PNA-I-treated animals were:

Exon10 forward primer: ggtcctcgtgtcacatctt

Exon9 reverse primer: cctcacctgatgcaagaaca

Exon8/10 reverse primer: tgcttctcggagccactc

PCR products were loaded onto 2% agarose gels (containing EtBr) along side a DNA mass

ruler. TBE running buffer was used for electrophoresis.

*Western blot*

Approximately 150 mg liver from mice treated with two different PNAs (PNA-I and PA5233,  
5 2x4.8  $\mu\text{mol/kg/day}$  for four days, PD-03-025) was homogenized in 4 ml detergent solution  
supplemented with proteinase inhibitors (Pro-Prep Solution from iNtRON Biotechnology).  
The samples were kept at  $-20^{\circ}\text{C}$  until analysis.

The total protein concentrations in liver samples were measured by using a detergent com-  
patible protein assay based on bicinchoninic acid (BCA) for calorimetric detection and quanti-  
10 fication (Pierce).

About 70  $\mu\text{g}$  total proteins from each sample was run on a 10% NuPage Bis-Tris gel (Invitro-  
gen) and blotted onto a PVDF membrane according to manufacturers instructions (Invitro-  
gen). The membrane was incubated with 0.25  $\mu\text{g/ml}$  anti-HNF4 $\alpha$  rabbit IgG targeting the C-  
terminal amino acid residues 455-465 of the human HNF4 $\alpha$  protein (Active Motif) followed by  
15 incubation with secondary antibodies and a chromogenic substrate from Invitrogen.


## 5 REFERENCES

1. Nielsen, P.E., Egholm, M., Berg, R.H. and Buchardt, O., *Science* (1991) **254**, 1497-1500.
2. Egholm, M., Buchardt, O., Christensen, L., Behrens, C., Freier, S. M., Driver, D.A., Berg, R.H., Kim, S.K., Norden, B., Nielsen, P.E., *Nature* (1993) **365**, 566-568.
- 10 3. Demidov, V., Potaman, V.N., Frank-Kamenetskii, M.D., Egholm, M., Buchardt, O. Sönnichsen, H. S., Nielsen, P.E., *Biochem. Pharmacol.* (1994) **48**, 1310-1313.
4. Nielsen, P.E. and Haaima, G., *Chemical Society Reviews* (1997) 73-78.
5. Hanvey J.C., Peffer N.J., Bisi J.E., Thomson S.A., Cadilla R., Josey J.A., Ricca D.J., Hassman C.F., Bonham M.A., Au K.G., *Science* (1992) **258** (5087), 1481-5.
- 15 6. Knudsen, H. and Nielsen, P.E., *Nucleic Acids Res.* (1996) **24**, 494-500.
7. Good, L. and Nielsen, P.E., *Proc. Natl. Acad. Sci. USA* (1998) **95**, 2073-2076.
8. Good, L. and Nielsen, P.E., *Nature Biotechnology* (1998) **16**, 355-358.
9. Lewis, L.G. et al. *Proc. Natl. Acad. Sci. USA* (1996) **93**, 3176-81.
10. Meyer, O. et al. *J. Biol. Chem.* (1998) **273**, 15621-7.
- 20 11. Nyce, J.W. and Metzger, W.J. *Nature* (1997) **385** 721-725.
12. Pooga, M. et al, *Nature Biotechnology* (1998) **16**, 857-61.
13. McMahon, B.M.; Mays, D.; Lipsky, J.; Stewart, J. A.; Fauq, A.; Richelson, E. *Antisense & Nucleic Acid Drug Development* **2002**, 12, 65-70.
14. Zhang, X.; Simmons, C. G.; Corey, D. R.; *Bioorganic & Medicinal Chemistry Letters*
- 25 **2001**, 11, 1269-1272.
15. Biessen, E. A. L.; Sliedregt-Bol, K.; Hoen, P.C.T.; Prince, P. Bilt, E. Valentijn, A. R. P. M.; Meeuwenoord, N. J.; Princen, H.; Bijsterbosch, M. K.; Marel, G. A., Boom, J. H., Berkel, T. J. C. *Bioconjugate Chem.* **2002**, 13, 295-302.
16. Litt. 5
- 30 17. Litt. 6
18. Litt. 1
19. Litt. 2
20. Litt. 3
21. Litt. 4
- 35 22. Litt. 7

23. Litt. 8
24. Litt. 9
25. Litt. 10
26. Litt. 11
- 5 27. Litt. 12
28. Litt. 13
29. Litt. 14
30. Litt. 15
31. Litt. 16
- 10 32. Ref.
33. Ref.
34. Litt. 17
35. Berge, S. M., Bighley L.D., Monkhouse D.C., *Pharmaceutical Science* (1977) 66, 1-19.

## CLAIMS


1. A modified Peptide Nucleic Acid (PNA) monomer of formula (I):



- 5 wherein B is a naturally-occurring nucleobase preferably A, T, G, or C, or a non-naturally-occurring nucleobase;

(Pr) is hydrogen or a protection group;


- 10  $R^1$ ,  $R^2$  and  $R^3$  are, independently, hydrogen, an amino acid side chain, or an  $C_{2-6}$ -alkyl, aryl, aralkyl, heteroaryl, hydroxy,  $C_{1-6}$ -alkoxy,  $C_{1-6}$ -alkylthio, hydroxy- or alkoxy- or alkylthio-substituted  $C_{1-6}$ -alkyl,  $-NR^4R^5$ , (wherein  $R^4$  and  $R^5$  independently are hydrogen,  $C_{1-6}$ -alkyl, hydroxy- or alkoxy- or alkylthio-substituted  $C_{1-6}$ -alkyl), or  $Z^1-Z^2$ , wherein  $Z^1$  is a bond or one of the radicals of formula (II):



- 15 wherein n is from 0 to 8;




and  $Z^2$  is  $\alpha$ - or  $\beta$ -forms of a monosaccharide, a disaccharide, a polysaccharide, or one of the radicals of formula (III):



5 provided that at least one of  $R^1$ ,  $R^2$ , or  $R^3$  is  $Z^1$ - $Z^2$ .

2. A compound according to claim 1, wherein the amino acid side chain is selected from the group consisting of  $C_{1-6}$ -alkyl, 3-guanidinopropyl, carboxymethyl, aminocarboxymethyl, mercaptomethyl, 2-carboxyethyl, aminocarboxyethyl, imidazol-4-yl-methyl, 4-aminobutyl, 2-(methylthio)ethyl, benzyl, hydroxymethyl, 1-hydroxyethyl, 3-indolyl, 4-hydroxybenzyl, 2-hydroxymethyl, or 3-ureidopropyl, 4-pyridomethyl (or 1-methylpropyl, 2-methylpropyl or 1-methylethyl).

3. A compound according to claim 1 or 2, wherein  $Z^1$  is one of the radicals of formula (IV) wherein  $n$  is from 0 to 8:




wherein  $n$  is from 0 to 8.

4. A compound according to claim 1, wherein the disaccharide is selected from the group consisting of  $\alpha$  and  $\beta$  forms of sucrose, lactose, and maltose.

5. A compound according to claim 1 or 3, wherein  $Z^1$ - $Z^2$  comprises ligands of formula

(V):



or beta-forms hereof.

5

6. A peptide nucleic acid oligomer with from 4 to 50 monomers selected from the group consisting of aeg-PNA monomers and at least one monomer of claim 1 to 5.

10

7. A peptide nucleic acid molecule comprising a peptide nucleic acid oligomer of claim 6 and a conjugate bound to said peptide nucleic acid either directly or through a linking moiety, wherein said conjugate is a reporter enzyme, a reporter molecule, a steroid, a carbohydrate, a terpene, a peptide, a protein, an aromatic lipophilic molecule, a non aromatic lipophilic molecule, a phospholipid, an intercalator, a cell receptor binding molecule, a crosslinking agent, a water soluble vitamin, a lipid soluble vitamin, an RNA/DNA cleaving complex, a metal chelator, a porphyrin, an alkylator, or a polymeric compound selected from polymeric amines, polymeric glycols and polyethers.

15

18

20

8. A peptide nucleic acid molecule comprising a peptide nucleic acid oligomer with from 4 to 50 aeg-PNA monomers and one or more ligands.


9. A peptide nucleic acid molecule of claim 8 wherein the ligand is one or more conjugate bound to said peptide nucleic acid either directly or through one or more linking

moieties, wherein said conjugate is a reporter enzyme, a reporter molecule, a steroid, a carbohydrate, a terpene, a peptide, a protein, an aromatic lipophilic molecule, a non aromatic lipophilic molecule, a phospholipid, an intercalator, a cell receptor binding molecule, a crosslinking agent, a water soluble vitamin, a lipid soluble vitamin, an RNA/DNA cleaving complex, a metal chelator, a porphyrin, an alkylator, or a polymeric compound selected from polymeric amines, polymeric glycols and polyethers.


10. A peptide nucleic acid molecule of claim 8 or 9 wherein the ligand has a high affinity towards a cancer related target.

11. A peptide nucleic acid molecule of claim 10 wherein the target is the  $\alpha v \beta 3$  integrin receptor.

12. A peptide nucleic acid molecule comprising a peptide nucleic acid oligomer with from 4 to 50 aeg-PNA monomers and one or more conjugates bound to said peptide nucleic acid either directly or through one or more linking moieties, wherein said conjugate is an amino acid side chain, or an  $C_{2-6}$ -alkyl, aryl, aralkyl, heteroaryl, hydroxy,  $C_{1-6}$ -alkoxy,  $C_{1-6}$ -alkylthio, hydroxy- or alkoxy- or alkylthio-substituted  $C_{1-6}$ -alkyl,  $-NR^4R^5$ , (wherein  $R^4$  and  $R^5$  independently are hydrogen,  $C_{1-6}$ -alkyl, hydroxy- or alkoxy- or alkylthio-substituted  $C_{1-6}$ -alkyl), or  $Z^1-Z^2$ , wherein  $Z^1$  is a bond, a peptide of from 1 to 10 amino acids or one of the radicals of formula (II):




(II)



wherein n is from 0 to 8;

and Z<sup>2</sup> is alfa- or beta forms of a monosaccharide, a disaccharide, a polysaccharide, or one of the radicals of formula (III):


5



10

13. A peptide nucleic acid molecule of claim 12 wherein the conjugate is a carbohydrate selected from the group consisting of  $\beta$ -D-galactosyl, 2-acetamido-2-deoxy-galactopyranosyl, 1-phenyl- $\beta$ -D-galactosyl, 1-propyl- $\beta$ -D-galactosyl or 1-butyl- $\beta$ -D-galactosyl

15



18


19

20


15. A peptide nucleic acid molecule of claim 13 or 14 wherein the carbohydrate is linked to the N-terminal end, the C-terminal end or to both ends of the peptide nucleic acid oligomer through one or more linkers.

16. A peptide nucleic acid molecule of the following formula:

5



17. A peptide nucleic acid molecule of the following formula:



10

18. A peptide nucleic acid molecule selected from:

(GalNAc(OH)<sub>3</sub>)<sub>2</sub>-Lys-Gly-CATCACTGGCAGACCCTG-NH<sub>2</sub>

(GalNAc(OH)<sub>3</sub>)<sub>3</sub>Lys<sub>2</sub>Gly-GTGGATGATACCTGGATC-NH<sub>2</sub>

15

or (GalNAc(OH)<sub>3</sub>)<sub>4</sub>Lys<sub>3</sub>Gly-GTGGATGATACCTGGATC-NH<sub>2</sub>

19. A peptide nucleic acid molecule according to any of the claims 9 to 15 wherein the linking moiety is an amino acid sequence of from 1 to 10 positively charged amino acids or amino acid analogues.

20

20. A peptide nucleic acid molecule according to claim 20 wherein the linking moiety is Lys-Lys-Lys-Lys.

21. Use of a PNA compound according to any of claims 1 to 20 in the manufacture of a medicament for the treatment or prevention of bacterial, viral, protozoal, and fungal infections, cancer, metabolic diseases, cardiovascular diseases, autoimmune and immunological disorders, or for disinfecting non-living objects, such as surgery tools, hospital inventory, dental tools, slaughterhouse inventory and tool, dairy inventory and tools, barbers and beauticians tools, and the like.
22. Use of a PNA compound according to any of claims 1 to 20 in the manufacture of a composition for the treatment or prevention of bacterial, viral, protozoal, and fungal infections, cancer, metabolic diseases, cardiovascular diseases, autoimmune and immunological disorders, or for disinfecting non-living objects, such as surgery tools, hospital inventory, dental tools, slaughterhouse inventory and tool, dairy inventory and tools, barbers and beauticians tools, and the like.
23. A pharmaceutical composition comprising, as an active ingredient, a compound according to any one of the preceding compound claims 1 to 20 or a pharmaceutically acceptable salt thereof together with a pharmaceutically acceptable carrier or diluent.
24. A composition according to claim 23 in unit dosage form, comprising from about 0.05 to about 100 mg, preferably from about 0.1 to about 50 mg of the compound according to any one of the preceding compound claims 1 to 20 or a pharmaceutically acceptable salt thereof.
25. A pharmaceutical composition according to any one of the claims 23 or 24 for oral, nasal, transdermal, pulmonal, or parenteral administration.
26. A pharmaceutical composition according to claim 23 to 25 for the treatment or prevention of bacterial, viral, protozoal, and fungal infections, cancer, metabolic diseases, cardiovascular diseases, autoimmune and immunological disorders, or treatment of non-living objects, the composition comprising, as an active ingredient, a compound according to any one of the preceding compound claims 1 to 20 or a pharmaceutically acceptable salt thereof together with a pharmaceutically acceptable carrier or diluent.

27. A method of treating a disease selected from bacterial, viral, protozoal, and fungal infections, cancer, metabolic diseases, cardiovascular diseases, autoimmune or immunological disorders comprising administering to a patient in need thereof an efficient amount of a compound of claim 1 to 20, the method comprising administering to a subject in need thereof an effective amount of a compound according to any one of the preceding compound claims 1 to 20 or a pharmaceutically acceptable salt thereof, or of a composition according to any one of the preceding composition claims.

28. The method according to claim 27, wherein the effective amount of the compound according to any one of the preceding compound claims 1 to 20 or a pharmaceutically acceptable salt or ester thereof is in the range of from about 0.05 to about 100 mg per day, preferably from about 0.1 to about 50 mg per day.

29. A compound selected from:

GalNAc(OBz)<sub>3</sub>-O-(CH<sub>2</sub>)<sub>4</sub>-COOH,

GalNAc(OBz)<sub>3</sub>-O-(CH<sub>2</sub>)<sub>4</sub>-CONH-Lys(GalNAc(OBz)<sub>3</sub>-O-(CH<sub>2</sub>)<sub>4</sub>-CONH-)-Gly-OH,

GalNAc(OBz)<sub>3</sub>-O-(CH<sub>2</sub>)<sub>4</sub>-CONH-Lys(GalNAc(OBz)<sub>3</sub>-O-(CH<sub>2</sub>)<sub>4</sub>-CONH)-

Lys(GalNAc(OBz)<sub>3</sub>-O-(CH<sub>2</sub>)<sub>4</sub>-CONH)-Gly-OH or

GalNAc(OBz)<sub>3</sub>-O-(CH<sub>2</sub>)<sub>4</sub>-CONH -Lys(GalNAc(OBz)<sub>3</sub>-O-(CH<sub>2</sub>)<sub>4</sub>-CONH -

Lys(GalNAc(OBz)<sub>3</sub>-O-(CH<sub>2</sub>)<sub>4</sub>-CONH -Lys(GalNAc(OBz)<sub>3</sub>-O-(CH<sub>2</sub>)<sub>4</sub>-CONH))) -Gly-OH

## ABSTRACT

The present invention concerns novel drugs for use in combating various diseases. More  
5 particular the invention concerns peptide nucleic acid (PNA) drugs, which are optionally  
modified in order to obtain novel PNA molecules with cell-specific delivery.






Figure 4